

Rec'd PCT/PTO 28 DEC 2004

#2

REC'D 11 AUG 2003

WIPO PCT

P1 1047285

THE UNITED STATES OF AMERICA

TO ALL TO WHOM THESE PRESENTS SHALL COME:

UNITED STATES DEPARTMENT OF COMMERCE
United States Patent and Trademark Office

August 05, 2003

THIS IS TO CERTIFY THAT ANNEXED HERETO IS A TRUE COPY FROM
THE RECORDS OF THE UNITED STATES PATENT AND TRADEMARK
OFFICE OF THOSE PAPERS OF THE BELOW IDENTIFIED PATENT
APPLICATION THAT MET THE REQUIREMENTS TO BE GRANTED A
FILING DATE.

APPLICATION NUMBER: 60/408,739

FILING DATE: September 06, 2002

RELATED PCT APPLICATION NUMBER: PCT/US03/20407

By Authority of the
COMMISSIONER OF PATENTS AND TRADEMARKS



H. Phillips
H. PHILLIPS
Certifying Officer


PRIORITY DOCUMENT
SUBMITTED OR TRANSMITTED IN
COMPLIANCE WITH
RULE 17.1(a) OR (b)

BEST AVAILABLE COPY

EXPRESS MAIL NO: EL889345381US
DATED: September 6, 2002

PROVISIONAL APPLICATION COVER SHEET

This is a request for filing a PROVISIONAL APPLICATION under 37 CFR 1.53(c).

DOCKET NUMBER		Docket No. PR1918-1		Type a plus sign (+) inside this box →	
INVENTOR(S)/APPLICANT(S)					
LAST NAME	FIRST NAME	MIDDLE INITIAL	RESIDENCE (CITY AND EITHER STATE OR FOREIGN COUNTRY)		
1) Desnoyers	Luc		San Francisco, CA		
APPLICATION IDENTIFIER (if no INVENTOR(S)/APPLICANT(S) are named)					
TITLE OF THE INVENTION (280 characters max)					
Methods and Compositions for Modulating and Detecting WISP Activity					
CORRESPONDENCE ADDRESS					
Diane L. Marschang  09157 PATENT TRADEMARK OFFICE			Telephone Number (650) 225-5416		
			Facsimile Number (650) 952-9881		
STATE	California	ZIP CODE	94080-4990	COUNTRY	USA
ENCLOSED APPLICATION PARTS (check all that apply)					
<input checked="" type="checkbox"/> Specification		Number of Pages <u>81</u> (including claims)			
<input checked="" type="checkbox"/> Drawings		Number of Pages <u>25</u>			
METHOD OF PAYMENT (check one)					
<input type="checkbox"/> A check or money order is enclosed to cover the Provisional filing fees.					
<input checked="" type="checkbox"/> The Commissioner is hereby authorized to charge filing fees and credit Deposit Account No. 07-0630. <u>A duplicate of this sheet is enclosed.</u>					
PROVISIONAL FILING FEE AMOUNT (\$) 160.00					

The invention was made by an agency of the United States Government or under a contract with an agency of the United States Government.

X No.

Respectfully submitted,
GENENTECH, INC.

By: Diane L. Marschang
Diane L. Marschang
Reg. No. 35,600

Date: September 6, 2002

Revised (12/1/97)

60428739.090602

EL889345381US: Express Mail Number

September 6, 2002: Date of Deposit

METHODS AND COMPOSITIONS FOR MODULATING AND DETECTING WISP ACTIVITY

Field of the Invention

5 The present invention relates generally to methods and compositions for use in modulating the activity(s) of WISP polypeptides, particularly, WISP-1 polypeptides. The invention also relates to methods and compositions for in vitro, in situ, and/or in vivo diagnosis and/or
10 treatment of mammalian cells or pathological conditions associated with WISP polypeptides.

Background of the Invention

15 Connective tissue growth factor (CTGF) is a growth factor induced in fibroblasts by many factors, including TGF- β , and is essential for the ability of TGF- β to induce anchorage-independent growth (AIG), a property of transformed cells. CTGF also is mitogenic and chemotactic for cells, and hence growth factors in this family are believed to play a role in the normal development, growth, and repair of human tissue. Five proteins
20 related to CTGF, including Cyr61, Nov, WISP-1, WISP2, and WISP-3 have been isolated, cloned, sequenced, and characterized as belonging to the CCN gene family. Oemar and Luescher, Arterioscler. Thromb. Vasc. Biol., 17: 1483-1489 (1997); Brigstock, Endocrine Rev., 20:189-206 (1999). The gene encoding Cyr61 has been found to promote angiogenesis, tumor growth, and
25 vascularization. Babic et al., Proc. Natl. Acad. Sci. USA, 95: 6355-6360 (1998). The nov gene is expressed in the kidney essentially at the embryonic stage, and alterations of nov expression, relative to the normal kidney, have been detected in both avian nephroblastomas and human Wilms' tumors. Martinerie et al., Oncogene, 9: 2729-2732 (1994). Wt1
30 downregulates human nov expression, which downregulation might represent a key element in normal and tumoral nephrogenesis. Martinerie et al., Oncogene, 12: 1479-1492 (1996).

The different members of the CCN family interact with various soluble or matrix associated macromolecules in particular sulfated glycoconjugates
35 (Bork, FEBS Letters, 327:125-130). This interaction was used to purify Cyr61 and CTGF by affinity chromatography on heparin-agarose (Frazier et al., J. Invest. Dermatol., 107:404-411 (1996); Kireeva et al., Mol. Cell. Biol., 16:1326-1334 (1996)). Cyr61 is secreted and associated with both the extracellular matrix and the cell surface due to its affinity for
40 heparan sulfate (Yang et al., Cell. Growth Diff., 2:351-357 (1991)). Recently, WISP-1 was shown to interact with decorin and biglycan, two secreted dermatan sulfate proteoglycans. (Desnoyers, et al., J. Biol. Chem., 276:47599-47607 (2001)).

CONFIDENTIAL

The murine protein ELM1 was recently identified in low metastatic cells. Hashimoto et al., J. Exp. Med., 187:289-296 (1998). The *elm1* gene, a mouse orthologue of WISP-1 disclosed below, is another member of the CNN gene family. It suppresses *in vivo* tumor growth and metastasis of K-1735 murine melanoma cells. Another recent article on rCop-1, the rat orthologue of WISP-2 described below, describes the loss of expression of this gene after cell transformation. Zhang et al., Mol. Cell. Biol., 18:6131-6141 (1998).

Wnts are encoded by a large gene family whose members have been found in round worms, insects, cartilaginous fish, and vertebrates. Holland et al., Dev. Suppl., 125-133 (1994). Wnts are thought to function in a variety of developmental and physiological processes since many diverse species have multiple conserved *Wnt* genes. McMahon, Trends Genet., 8: 236-242 (1992); Nusse and Varmus, Cell, 69: 1073-1087 (1992). *Wnt* genes encode secreted glycoproteins that are thought to function as paracrine or autocrine signals active in several primitive cell types. McMahon, *supra* (1992); Nusse and Varmus, *supra* (1992). The *Wnt* growth factor family includes more than ten genes identified in the mouse (*Wnt*-1, -2, -3A, -3B, -4, -5A, -5B, -6, -7A, -7B, -8A, -8B, -10B, -11, -12, and -13) (see, e.g., Gavin et al., Genes Dev., 4: 2319-2332 (1990); Lee et al., Proc. Natl. Acad. Sci. USA, 92: 2268-2272 (1995); Christiansen et al., Mech. Dev., 51: 341-350 (1995)) and at least nine genes identified in the human (*Wnt*-1, -2, -3, -5A, -7A, -7B, -8B, -10B, and -11) by cDNA cloning. See, e.g., Vant Veer et al., Mol. Cell. Biol., 4: 2532-2534 (1984).

The *Wnt*-1 proto-oncogene (*int*-1) was originally identified from mammary tumors induced by mouse mammary tumor virus (MMTV) due to an insertion of viral DNA sequence. Nusse and Varmus, Cell, 31: 99-109 (1982). In adult mice, the expression level of *Wnt*-1 mRNA is detected only in the testis during later stages of sperm development. *Wnt*-1 protein is about 42 KDa and contains an amino-terminal hydrophobic region, which may function as a signal sequence for secretion (Nusse and Varmus, *supra*, 1992). The expression of *Wnt*-2 is detected in mouse fetal and adult tissues and its distribution does not overlap with the expression pattern for *Wnt*-1. *Wnt*-3 is associated with mouse mammary tumorigenesis. The expression of *Wnt*-3 in mouse embryos is detected in the neural tubes and in the limb buds. *Wnt*-5a transcripts are detected in the developing fore- and hind limbs at 9.5 through 14.5 days and highest levels are concentrated in apical ectoderm at the distal tip of limbs. Nusse and Varmus, *supra* (1992). Recently, a *Wnt* growth factor, termed *Wnt*-x, was described (WO95/17416) along with the detection of *Wnt*-x expression in bone tissues and in bone-derived cells. Also described was the role of *Wnt*-x in the maintenance of mature osteoblasts and the use of the *Wnt*-x growth factor as

therapeutic agent or in the development of other therapeutic agents to treat bone-related diseases.

Wnts may play a role in local cell signaling. Peifer and Polakis, Science, 287:1606-1609 (2000). Biochemical studies have shown that much of the secreted Wnt protein can be found associated with the cell surface or extracellular matrix rather than freely diffusible in the medium. Papkoff and Schryver, Mol. Cell. Biol., 10: 2723-2730 (1990); Bradley and Brown, EMBO J., 9: 1569-1575 (1990).

Studies of mutations in *Wnt* genes have indicated a role for Wnts in growth control and tissue patterning. In *Drosophila*, *wingless* (*wg*) encodes a *Wnt*-related gene (Rijsewijk et al., Cell, 50: 649-657 (1987)) and *wg* mutations alter the pattern of embryonic ectoderm, neurogenesis, and imaginal disc outgrowth. Morata and Lawrence, Dev. Biol., 56: 227-240 (1977); Baker, Dev. Biol., 125: 96-108 (1988); Klingensmith and Nusse, Dev. Biol., 166: 396-414 (1994). In *Caenorhabditis elegans*, *lin-44* encodes a Wnt homolog which is required for asymmetric cell divisions. Herman and Horvitz, Development, 120: 1035-1047 (1994). Knock-out mutations in mice have shown Wnts to be essential for brain development (McMahon and Bradley, Cell, 62: 1073-1085 (1990); Thomas and Cappechi, Nature, 346: 847-850 (1990)), and the outgrowth of embryonic primordia for kidney (Stark et al., Nature, 372: 679-683 (1994)), tail bud (Takada et al., Genes Dev., 8: 174-189 (1994)), and limb bud. Parr and McMahon, Nature, 374: 350-353 (1995). Overexpression of *Wnt-1* in the mammary gland can result in mammary hyperplasia (McMahon, *supra* (1992); Nusse and Varmus, *supra* (1992)), precocious alveolar development (Bradbury et al., Dev. Biol., 170: 553-563 (1995)), and mammary adenocarcinomas (Li et al., Oncogene, 19:1002-1009 (2000)).

Wnt-5a and *Wnt-5b* are expressed in the posterior and lateral mesoderm and the extraembryonic mesoderm of the day 7-8 murine embryo. Gavin et al., *supra* (1990). These embryonic domains contribute to the AGM region and yolk sac tissues from which multipotent hematopoietic precursors and HSCs are derived. Dzierzak and Medvinsky, Trends Genet., 11: 359-366 (1995); Zon et al., in Gluckman and Coulombel, ed., Colloque, INSERM, 235: 17-22 (1995), presented at the Joint International Workshop on Foetal and Neonatal Hematopoiesis and Mechanism of Bone Marrow Failure, Paris France, April 3-6, 1995; Kanatsu and Nishikawa, Development, 122: 823-830 (1996). *Wnt-5a*, *Wnt-10b*, and other Wnts have been detected in limb buds, indicating possible roles in the development and patterning of the early bone microenvironment as shown for *Wnt-7b*. Gavin et al., *supra* (1990); Christiansen et al., Mech. Devel., 51: 341-350 (1995); Parr and McMahon, *supra* (1995).

For a review on Wnt, see Cadigan and Nusse, Genes & Dev., 11: 3286-3305 (1997).

Pennica et al., Proc. Natl. Acad. Sci., 95:14717-14722 (1998) describe the cloning and characterization of two genes, *WISP-1* and *WISP-2*, that, are up-regulated in the mouse mammary epithelial cell line C57MG transformed by Wnt-1, and a third related gene, *WISP-3*. (See also, WO 99/21998 published May 6, 1999; WO 99/21999 published May 6, 1999). Pennica et al. report that these *WISP* genes may be downstream of Wnt-1 signaling and that aberrant levels of *WISP* expression in colon cancer may play a role in colon tumorigenesis. *WISP-1* has recently been identified as a β -catenin-regulated gene and the characterization of its oncogenic activity demonstrated that *WISP-1* might contribute to β -catenin-mediated tumorigenesis (Xu et al., Gene & Develop., 14:585-595 (2000)). *WISP-1* overexpression in normal rat kidney cells (NRK-49F) induced morphological transformation, accelerated cell growth and enhanced saturation density. In addition, these cells readily form tumors when injected into nude mice suggesting that *WISP-1* may play some role in tumorigenesis (Xu et al., supra 2000). *WISP-1* is also overexpressed in transformed human breast cancer cell lines and in about 47% of primary human breast cancer associated with certain advanced features. Xie et al., Cancer Res., 61:8917-8923 (2001); Saxena et al., Mol. Cell Biochem., 228:99-104 (2001); Michaelson et al., Oncogene, 20:5093-5099 (2001). A particular *WISP-1* variant has also been reported to be overexpressed in about 86% of human scirrhous gastric carcinoma cells. Tanaka et al., Oncogene, 20:5525-5532 (2001).

Hurvitz et al., Nature Genetics, 23:94-97 (1999) describe a study involving *WISP3* in which nine different mutations of *WISP3* in unrelated individuals were found to be associated with the autosomal recessive skeletal disorder, progressive pseudorheumatoid dysplasia (PPD). *WISP3* expression by RT-PCR was observed by Hurvitz et al. in human synoviocytes, articular cartilage chondrocytes, and bone-marrow-derived mesenchymal progenitor cells.

PCT application WO98/21236 published May 22, 1998 discloses a human connective tissue growth factor gene-3 (CTGF-3) encoding a 26 kD member of the growth factor superfamily. WO98/21236 discloses that the CTGF-3 amino acid sequence was deduced from a human osteoblast cDNA clone, and that CTGF-3 was expressed in multiple tissues like ovary, testis, heart, lung, skeletal muscle, adrenal medulla, adrenal cortex, thymus, prostate, small intestine and colon.

Hyaluronic acid (also referred to as HA, hyaluronate, or hyaluronan) is recognized in the literature as being an important component of the extracellular matrix (See, e.g., Hardingham et al., FASEB J., 6:861-870

1992); Laurent et al., FASEB J., 6:2397-2404 (1992)). HA is a component of skin and mesenchymal tissues where it facilitates cell migration during wound healing, inflammation, and embryonic morphogenesis. (Knudson et al., FASEB J., 7:1233-1241 (1993); Knudson et al., CIBA Found. Symp., 143:150-169 (1989)). HA has also been reported to play a role in certain types of metastases. (Naor et al., CD44: Structure, Function and Association with the Malignant Process, Advances in Cancer Research, Academic Press (1997), pages 241-319). The largest concentrations of HA are found in the skin and musculo-skeletal system which account for over 50% of total body HA. (Banerji et al., J. Cell Biol., 144:789-801 (1999)).

Various investigators have reported on receptors which bind HA. One of the receptors identified for HA is the CD44 protein. (See, e.g., Culty et al., J. Cell Biology, 111:2765-2774 (1990); Aruffo et al., Cell, 61:1303-1313 (1990); Naor et al., CD44: Structure, Function and Association with the Malignant Process, Advances in Cancer Research, Academic Press (1997), pages 241-319); Ropponen et al., Cancer Res., 58:342-347 (1998); Masaki et al., Cancer, 92:2539-2546 (2001). CD44 is a family of cell-surface glycoproteins generated from a single gene by alternative splicing and differential glycosylation. (Wielenga et al., Am. J. Pathology, 154:515-523 (1999)). CD44 is believed to function as a cell adhesion receptor, linking extracellular molecules, specifically hyaluronate, to the cell and the cytoskeleton (Wielenga et al., supra). CD44 is expressed on epithelial, mesenchymal and lymphoid cells. (Lesley et al., Adv. Immunol., 54:271-335 (1994)). Wielenga et al. report that CD44 expression may be regulated by the WNT pathway, based on certain experiments analyzing CD44 expression in the intestinal mucosa of mice and humans with genetic defects in either APC or Tcf-4. (Wielenga et al., supra).

Other HA receptors characterized to date include RHAMM (also referred to as receptor for hyaluronic acid mediated motility), a 58 kD intracellular protein expressed transiently on the surface of transformed lymphocytes (Hardwick et al., J. Cell Biol., 117:1343-1350 (1992); Turley et al., Exp. Cell Res., 207:277-282 (1993)). RHAMM expression in fibroblasts was reported to promote metastasis and play an important role in H-Ras transformation (Hall et al., infra).

Another receptor which binds HA was described by Banerji et al. (Banerji et al., supra). Banerji et al. report a receptor on lymph vessel walls, referred to as "LYVE-1", which is a 322-residue type I integral membrane polypeptide which has a 41% similarity to the CD44 receptor. Unlike the CD44 receptor for HA, the LYVE-1 protein is absent in blood vessels. In addition, layilin (Bono et al., Mol. Biol. Cell, 12:891-900 (2001)) and HARE (Zhou et al., J. Biol. Chem., 275:37733-37741 (2000)) were also described as HA receptors.

Summary of the Invention

Applicant has surprisingly found that WISP-1 can induce HAS2 (hyaluronan synthase 2), CD44, and RHAMM mRNA expression, CD44 protein synthesis, and HA secretion. The induction or secretion of such molecules may promote or increase cancer cell growth, motility and/or metastatic potential. The present invention thus provides for instance, WISP-1 antagonists and methods of using such antagonists. The antagonists described herein find utility for, among other things, *in vitro*, *in situ*, or *in vivo* diagnosis or treatment of mammalian cancer cells or other pathological conditions associated with the induction or secretion of HAS2, HA, CD44 or RHAMM.

In embodiments of the invention, there are provided isolated WISP-1 antagonists. Such antagonists may comprise antibodies, such as WISP-1 antibodies. In preferred embodiments, the antagonists may block or neutralize WISP-1 induction or secretion of HAS2, HA, CD44 or RHAMM. Such antagonistic antibodies may, for example, be monoclonal antibodies, chimeric antibodies, humanized antibodies, or human antibodies. The WISP-1 antagonists contemplated for use in the invention include WISP-1 immunoadhesins, WISP-1 variants, covalently modified forms thereof, or fusion proteins thereof. By way of example, such antagonists may include pegylated WISP-1 or WISP-1 fused to heterologous sequences such as epitope tags or leucine zippers. The methods contemplate the use of a single type of antagonist molecule or a combination of two or more types of antagonists.

The methods of the invention include methods to treat pathological conditions or diseases in mammals associated with or resulting from WISP-1, including, induction or secretion of HAS2, HA, CD44 or RHAMM by WISP-1. In the methods of treatment, WISP-1 antagonists may be administered to the mammal suffering from such pathological condition or disease. For example, the invention provides a method comprising exposing a mammalian cell(s), such as a cancer cell(s), to one or more WISP-1 antagonists in an amount effective to decrease, neutralize or block WISP-1 induction or secretion of HAS2, HA, CD44 or RHAMM. The cell may be in cell culture or in a mammal, e.g. a mammal suffering from, for instance, cancer.

The invention also provides compositions which comprise one or more WISP-1 antagonists. Optionally, the compositions of the invention will include pharmaceutically acceptable carriers or diluents. Preferably, the compositions will include one or more WISP-1 antagonists in an amount which is therapeutically effective to treat a pathological condition or disease.

The invention also provides articles of manufacture and kits which include one or more WISP-1 antagonists.

The invention also provides methods of conducting screening assays to identify candidate molecules, such as small molecule compounds, polypeptides or antibodies, which act as antagonists with respect to blocking or neutralizing WISP-1 induction or secretion of HAS2, HA, CD44 or RHAMM.

More particular embodiments of the invention include isolated WISP-1 antagonists which inhibit or neutralize the induction or secretion of HAS2, HA, CD44 or RHAMM by native WISP-1 polypeptide in at least one type of mammalian cell, said antagonist being selected from the group consisting of an anti-WISP-1 antibody, a WISP-1 immunoadhesin, a WISP-1 variant, and fusion proteins thereof. The antagonist may comprise an anti-WISP-1 antibody which binds native human WISP-1 polypeptide comprising amino acids 23-367 of Figures 9A-9C or one or more domains of WISP-1 comprising amino acids encoded by the sequences of SEQ ID NO:3; 4; 5; 6; 7; 8; 9; 10; or 11 herein.

The invention also provides compositions comprising the antagonists described herein and a carrier, optionally the carrier is a pharmaceutically-acceptable carrier.

The invention also provides methods of inhibiting or neutralizing WISP-1 induction or secretion of HAS2, HA, CD44 or RHAMM in mammalian cells, comprising exposing said mammalian cells to an effective amount of WISP-1 antagonist, wherein said WISP-1 antagonist is selected from the group consisting of

- a) a WISP-1 immunoadhesin;
- b) a WISP-1 polypeptide linked to a nonproteinaceous polymer selected from the group consisting of polyethylene glycol, polypropylene glycol, and polyoxyalkylene;
- c) a WISP-1 antibody; and
- d) a WISP-1 variant.

The WISP-1 immunoadhesins employed in the methods may comprise a WISP-1 sequence fused to a Fc region of an immunoglobulin. The anti-WISP-1 antibodies employed in the methods may bind native human WISP-1 comprising amino acids 23-367 of Figures 9A-9C, or one or more of the domains of WISP-1 described in the Examples below. In the methods, the mammalian cells may comprise cancer cells, and optionally the cells comprise colon or colorectal cancer cells, breast cancer cells, lung cancer cells, or brain cancer cells (such as glioma or glioblastoma).

In other embodiments, there are provided methods of treating cancer in a mammal, comprising administering to said mammal an effective amount of WISP-1 antagonist. Optionally, in said methods, the antagonist may inhibit

neutralize induction or secretion of HAS2, HA, CD44 or RHAMM by native human WISP-1 polypeptide in at least one type of mammalian cell and said antagonist is selected from the group consisting of an anti-WISP-1 antibody, a WISP-1 immunoadhesin and a WISP-1 variant. Optionally, the antagonist may act to inhibit cancer cell growth or cancer cell metastasis. The cancer in the mammal may comprise colon or colorectal cancer cells, breast cancer cells, lung cancer cells, or brain cancer cells (such as glioma or glioblastoma). Optionally, the antagonist(s) employed in the methods inhibits or reduces cancer cell growth or metastasis. Also, in the methods, chemotherapy or radiation may also be administered to the mammal.

In more particular embodiments, there are provided antibodies which specifically bind to one or more domains of WISP-1 polypeptide (described further in the Examples below) comprising amino acids encoded by the sequences of SEQ ID NO:3; 4; 5; 6; 7; 8; 9; 10; or 11 herein. Optionally, the antibody is a monoclonal antibody. Optionally, the monoclonal antibody comprises the 3D11, 11C2, 9C10, 5D4, or 9C11 antibody secreted by the hybridoma deposited with ATCC as accession number __, __, __, __, or __, respectively.

Also provided are antibodies which bind to the same epitope as the epitope to which the 3D11, 11C2, 9C10, 5D4, or 9C11 monoclonal antibody produced by the hybridoma cell line deposited as ATCC accession number __, __, __, __, or __, respectively, binds.

In yet other particular embodiments, there is provided the hybridoma cell line which produces monoclonal antibody 3D11, 11C2, 9C10, 5D4, or 9C11 and deposited with ATCC as accession number __, __, __, __, or __, respectively, and the monoclonal antibody 3D11, 11C2, 9C10, 5D4, or 9C11 secreted by the hybridoma deposited with ATCC as accession number __, __, __, __, or __, respectively.

There are also provided isolated anti-WISP-1 monoclonal antibodies, comprising antibodies which bind to WISP-1 polypeptide and competitively inhibit binding of the monoclonal antibody produced by the hybridoma deposited as ATCC accession no. __, __, __, __, or __ to said WISP-1 polypeptide.

There are also provided chimeric or humanized anti-WISP-1 antibodies which specifically bind to WISP-1 polypeptide and comprise (a) a sequence derived from the 3D11, 11C2, 9C10, 5D4, or 9C11 antibody secreted by the hybridoma deposited with ATCC as accession number __, __, __, __, or __, respectively. Optionally, such antibodies may comprise a heavy chain, light chain or variable regions derived from the 3D11, 11C2, 9C10, 5D4, or 9C11 antibody.

The anti-WISP-1 antibodies may be linked to one or more non-proteinaceous polymers selected from the group consisting of polyethylene

Brief Description of the Drawings

Figures 1A-1F show WISP-1 promotion of hyaluronan production. Particle exclusion assay demonstrating a hyaluronan coat at the surface of NRK/WISP-1H (A) and NRK/WISP-1L (B) but absent from NRK/control (C). NRK/WISP-1H (D) and NRK/control (E) were stained for hyaluronan with bHABP. F, Time course of HA accumulation in NRK/WISP-1H and NRK/control media.

Figures 2A-2C show WISP-1 increases HAS2, CD44 and RHAMM mRNA expression and CD44 protein expression. A, Real time RT-PCR analysis of HAS1, HAS2, HAS3, CD44, RHAMM and hyaluronidase (Hyal) mRNA expression in NRK/WISP-1H, NRK/WISP-1L and NRK/control cells. Results are given as fold induction over the expression in NRK/control cells. B, Flow cytometry analysis of CD44 expression in NRK/WISP-1H and NRK/control cells. The shaded area represents the fluorescence intensity of the secondary antibody alone. C, Western blot analysis of CD44 protein in NRK/WISP-1H and NRK/control cells. Actin staining was used as a loading control.

Figures 3A-3G show WISP-1 expression increases cell motility and modifies cellular morphology. NRK/control cells (A) formed defined colonies when plated at low density whereas NRK/WISP-1L (B) and NRK/WISP-1H cells (C) scattered. NRK/WISP-1H (E) showed a de-differentiated spindloid morphology with lamellipodia compared to NRK/control cells (D). F, The random migration of NRK/WISP-1H and NRK/control cells was measured over 15 hours using time lapse microscopy. Results represent the typical average migration distance of cells in one field. G, Motility of NRK/WISP-1H and NRK/control cells was evaluated by a cell wound healing assay and measured after 15 hours using time lapse microscopy. Data represent the results of a typical experiment.

Figures 4A-4C show WISP-1 addition induces HAS2 mRNA expression and haptotactic migration of NRK cells. A, Real time RT-PCR analysis of HAS2 and CD44 expression in NRK cells seeded on coated surface. In certain cases the coated substrates were further incubated with WISP-1 (treatment). Results are given in fold induction over expression in NRK cells plated on non-coated, non-treated surface. The underside of modified Boyden chamber filters were coated and NRK cells (B) or SW480 cells (C) were added to the upper chamber. In certain cases, additions were made directly to the lower chamber. Cell motility was evaluated by counting the cells that migrated to the lower side of the insert.

Figures 5A-5E show WISP-1, HAS2 and CD44 over-expression and increased cell motility in C57MG/Wnt-1 cells and MMTV-Wnt-1 transgenic mice mammary tumors. A, Real time RT-PCR analysis of WISP-1, HAS2 and CD44 expression in C57MG/Wnt-1 and C57MG/control cells. Results are given as 5 fold induction over expression in C57MG/control cells. B, Western blot analysis of CD44 content in C57MG/Wnt-1 and C57MG/control cells. Actin staining was used as a loading control. C, C57MG/control cells formed distinct colonies when plated at low density whereas C57MG/Wnt-1 cells scattered. Semi-quantitative RT-PCR analysis of HAS2 (D) and CD44 (E) in 10 mammary tumors of MMTV-Wnt-1 transgenic mice. Results are expressed as relative fold induction over the expression in normal mammary gland tissue.

Figures 6A-6H show WISP-1, HAS2 and CD44 expression and HA accumulation in MMTV-Wnt-1 transgenic mammary tumors. In situ hybridization of WISP-1 (A, B) HAS2 (C, D) and CD44 (E, F), 15 immunohistochemistry of CD44 (G) and bHABP fluorescent staining of hyaluronan (H) in MMTV-Wnt-1 transgenic mouse mammary tumors. t, tumor; s, stroma.

Figures 7A-7L show WISP-1 expression promotes metastatic lung colonization and tumor growth. The effect of WISP-1 was analyzed by 20 injecting NRK/control, NRK/WISP-1L or NRK/WISP-1H cells in the tail vein of nude mice. At various times after injection, lungs were imaged by MRI (A, D, G, J), excised and their gross appearance (B, E, H, K) and histological features (C, F, I, L) were recorded. The severity of lung invasion by tumors was graded as normal (A, B, C); grade I (D, E, F); grade II (G, H, 25 I) or grade III (J, K, L).

Figures 8A-B show CD44 antibody prevents NRK/WISP-1H cell metastasis. NRK/WISP-1H cells (2.5×10^5 cells) were injected in the tail vein of nude mice. Mice were treated twice a week with 10 mg/kg of a CD44 antibody, an isotype control antibody or with buffer only (PBS). The lungs were excised 30 after four weeks for gross anatomical analysis. Picture of a normal lung is shown for comparison.

Figures 9A-9C show the nucleotide (SEQ ID NO:2) and putative amino acid sequence (SEQ ID NO:1) of native human WISP-1.

Figure 10 shows the effects of WISP-1 expression on NRK metastatic 35 potential.

Figures 11A-B show the binding properties of WISP-1 antibodies to various WISP-1 constructs and preparations.

Figures 12A-G show diagrams of various domains of WISP-1 polypeptide and the results of assays conducted to identify the epitopes recognized by 40 the WISP-1 antibodies 11C2, 5D4, 9C11, and 3D11, respectively.

Figure 13 shows the results on an assay conducted to identify the epitope recognized by the WISP-1 antibody 9C10.

Figure 14 shows the results of an ELISA demonstrating detection of WISP-1 by WISP-1 antibodies.

Figure 15 shows the results of an ELISA binding assay demonstrating that WISP-1 antibodies recognizing the variable region of WISP-1 can inhibit WISP-1 binding to heparin.

Figures 16A-B show the results of an assay detecting inhibition of haptotaxis of NRK cells by WISP-1 antibodies (16A) and a chart summarizing the properties and characteristics of WISP-1 antibodies (16B).

Figures 17A-E show the results of an in vivo study of the effects of WISP-1 antibodies. After 3 weeks, the severity of the lesions found in WISP-1 antibody treated animals was greatly attenuated compared to control (Fig. 17 a, b). The number of nodules and the average area of the metastatic foci found in mice treated with WISP-1 antibodies (n=5) were reduced compared to animals treated with a control antibody (Fig. 17 c,d). The total pulmonary area covered by the lesions was reduced by 82-97% compared to animal treated with an isotype control antibody (Fig. 17e)

Detailed Description of the Invention

I. Definitions

The term "WISP polypeptide" refers to the family of native-sequence human and mouse WISP proteins and variants described herein whose genes are induced at least by Wnt-1. This term includes WISP-1 and variants thereof.

Such WISP-1 proteins are described further below and in PCT application WO99/21998 published May 6, 1999, US 6,387,657 B1 issued May 14, 2002, and in Pennica et al., Proc. Natl. Acad. Sci., 95:14717-14722 (1998).

The terms "WISP-1 polypeptide", "WISP-1 homologue" and grammatical variants thereof, as used herein, encompass native-sequence WISP-1 protein and variants (which are further defined herein). The WISP-1 polypeptide may be isolated from a variety of sources, such as from human tissue types or from another source, or prepared by recombinant or synthetic methods, or by any combination of these and similar techniques.

The term "native-sequence WISP-1 polypeptide" comprises a polypeptide having the same amino acid sequence as a WISP-1 polypeptide derived from nature. Such native-sequence WISP-1 polypeptides can be isolated from nature or can be produced by recombinant or synthetic means. The term "native-sequence WISP-1 polypeptide" specifically encompasses naturally occurring truncated or secreted forms of a WISP-1 polypeptide disclosed herein, naturally occurring variant forms (e.g., alternatively spliced forms or splice variants), and naturally occurring allelic variants of a WISP-1 polypeptide. In one embodiment of the invention, the native-sequence WISP-1 polypeptide is a mature or full-length native-sequence human WISP-1 polypeptide comprising amino acids 23 to 367 of Figure 9 or

amino acids 1 to 367 of Figure 9, respectively, with or without N-terminal methionine.

In another embodiment of the invention, the native-sequence WISP-1 polypeptide is the full-length or mature native-sequence human WISP-1 polypeptide comprising amino acids 23 to 367 or 1 to 367 of Figure 9 wherein the valine residue at position 184 or the alanine residue at position 202 has/have been changed to an isoleucine or serine residue, respectively, with or without N-terminal methionine. In another embodiment of the invention, the native-sequence WISP-1 polypeptide is the full-length or, mature native-sequence human WISP-1 polypeptide comprising amino acids 23 to 367 or 1 to 367 of the Figure 9 wherein the valine residue at position 184 and the alanine residue at position 202 has/have been changed to an isoleucine or serine residue, respectively, with or without N-terminal methionine.

In another embodiment of the invention, the native-sequence WISP-1 polypeptide is one which is encoded by a nucleotide sequence comprising one of the human WISP-1 splice or other native-sequence variants, including SEQ ID NOS:23, 24, 25, 26, 27, 28, or 29 shown in WO99/21998, with or without an N-terminal methionine.

The term "WISP-1 variant" means an active WISP-1 polypeptide as defined below having at least about 80%, preferably at least about 85%, 86%, 87%, 88%, 89%, more preferably at least about 90%, 91%, 92%, 93%, 94%, most preferably at least about 95%, 96%, 97%, 98%, or 99% amino acid sequence identity with human mature WISP-1 having the deduced amino acid sequence shown in Fig. 9, or a soluble fragment thereof. Such variants include, for instance, WISP-1 polypeptides wherein one or more amino acid residues are added to, or deleted from, the N- or C-terminus of the full-length or mature sequences of Figure 9, or WISP-1 polypeptides wherein one or more amino acid residues are inserted or deleted from the internal sequence or domains of the polypeptide, including variants from other species, but excludes a native-sequence WISP-1 polypeptide. Preferably such a variant acts as an antagonist, as defined below.

An "extracellular domain", "ECD" or "secreted" protein refers to a form of a polypeptide which is essentially free of any transmembrane and cytoplasmic domains. A "secreted" form of a protein or polypeptide may or may not include a signal sequence.

"Stringent conditions" are those that (1) employ low ionic strength and high temperature for washing, for example, 0.015 M sodium chloride/0.0015 M sodium citrate/0.1% sodium dodecyl sulfate at 50°C; (2) employ during hybridization a denaturing agent, such as formamide, for example, 50% (vol/vol) formamide with 0.1% bovine serum albumin/0.1% Ficoll/0.1% polyvinylpyrrolidone/50 mM sodium phosphate buffer at pH 6.5

with 750 mM sodium chloride, 75 mM sodium citrate at 42°C; (3) employ 50% formamide, 5 x SSC (0.75 M NaCl, 0.075 M sodium citrate), 50 mM sodium phosphate (pH 6.8), 0.1% sodium pyrophosphate, 5 x Denhardt's solution, sonicated salmon sperm DNA (50 µg/ml), 0.1% SDS, and 10% dextran sulfate at 42°C, with washes at 42°C in 0.2 x SSC and 0.1% SDS; or (4) employ a buffer of 10% dextran sulfate, 2 x SSC (sodium chloride/sodium citrate), and 50% formamide at 55°C, followed by a high-stringency wash consisting of 0.1 x SSC containing EDTA at 55°C.

"Moderately stringent conditions" are described in Sambrook et al., Molecular Cloning: A Laboratory Manual (New York: Cold Spring Harbor Laboratory Press, 1989), and include the use of a washing solution and hybridization conditions (e.g., temperature, ionic strength, and percent SDS) less stringent than described above. An example of moderately stringent conditions is a condition such as overnight incubation at 37°C in a solution comprising: 20% formamide, 5 x SSC (150 mM NaCl, 15 mM trisodium citrate), 50 mM sodium phosphate (pH 7.6), 5 x Denhardt's solution, 10% dextran sulfate, and 20 mg/mL denatured sheared salmon sperm DNA, followed by washing the filters in 1 x SSC at about 37-50°C. The skilled artisan will recognize how to adjust the temperature, ionic strength, etc., as necessary to accommodate factors such as probe length and the like.

"Isolated," when used to describe the various polypeptides disclosed herein, means polypeptide that has been identified and separated and/or recovered from a component of its natural environment. Contaminant components of its natural environment are materials that would typically interfere with diagnostic or therapeutic uses for the polypeptide, and may include enzymes, hormones, and other proteinaceous or non-proteinaceous solutes. In preferred embodiments, the polypeptide will be purified (1) to a degree sufficient to obtain at least 15 residues of N-terminal or internal amino acid sequence by use of a spinning cup sequenator, or (2) to homogeneity by SDS-PAGE under non-reducing or reducing conditions using Coomassie blue or, preferably, silver stain. Isolated polypeptide includes polypeptide *in situ* within recombinant cells, since at least one component of the WISP natural environment will not be present. Ordinarily, however, isolated polypeptide will be prepared by at least one purification step.

The term "control sequences" refers to DNA sequences necessary for the expression of an operably linked coding sequence in a particular host organism. The control sequences that are suitable for prokaryotes, for example, include a promoter, optionally an operator sequence, and a ribosome binding site. Eukaryotic cells are known to utilize promoters, polyadenylation signals, and enhancers.

Nucleic acid is "operably linked" when it is placed into a functional relationship with another nucleic acid sequence. For example, DNA for a

presequence or secretory leader is operably linked to DNA for a polypeptide if it is expressed as a preprotein that participates in the secretion of the polypeptide; a promoter or enhancer is operably linked to a coding sequence if it affects the transcription of the sequence; or a ribosome binding site is operably linked to a coding sequence if it is positioned so as to facilitate translation. Generally, "operably linked" means that the DNA sequences being linked are contiguous, and, in the case of a secretory leader, contiguous and in reading phase. However, enhancers do not have to be contiguous. Linking is accomplished by ligation at convenient restriction sites. If such sites do not exist, the synthetic oligonucleotide adaptors or linkers are used in accordance with conventional practice.

A "liposome" is a small vesicle composed of various types of lipids, phospholipids and/or surfactant which is useful for delivery of a drug (such as the insulin and insulin variants disclosed herein) to a mammal. The components of the liposome are commonly arranged in a bilayer formation, similar to the lipid arrangement of biological membranes.

The terms "amino acid" and "amino acids" refer to all naturally occurring L-alpha-amino acids. This definition is meant to include norleucine, ornithine, and homocysteine. The amino acids are identified by either the single-letter or three-letter designations:

Asp	D	aspartic acid	Ile	I	isoleucine
Thr	T	threonine	Leu	L	leucine
Ser	S	serine	Tyr	Y	tyrosine
25 Glu	E	glutamic acid	Phe	F	phenylalanine
Pro	P	proline	His	H	histidine
Gly	G	glycine	Lys	K	lysine
Ala	A	alanine	Arg	R	arginine
Cys	C	cysteine	Trp	W	tryptophan
30 Val	V	valine	Gln	Q	glutamine
Met	M	methionine	Asn	N	asparagine

In the Sequence Listing and Figures, certain other single-letter or three-letter designations may be employed to refer to and identify two or more amino acids or nucleotides at a given position in the sequence.

"Percent (%) amino acid sequence identity" with respect to the polypeptide sequences identified herein is defined as the percentage of amino acid residues in a candidate sequence that are identical with the amino acid residues in such a polypeptide sequence identified herein, after aligning the sequences and introducing gaps, if necessary, to achieve the maximum percent sequence identity, and not considering any conservative substitutions as part of the sequence identity. Alignment for purposes of

determining percent amino acid sequence identity can be achieved in various ways that are within the skill in the art, for instance, using publicly available computer software such as BLAST, BLAST-2, ALIGN, ALIGN-2 or Megalign (DNASTAR) software. Those skilled in the art can determine appropriate parameters for measuring alignment, including any algorithms needed to achieve maximal alignment over the full-length of the sequences being compared. For purposes herein, however, % amino acid sequence identity values are obtained as described below by using the sequence comparison computer program ALIGN-2, wherein the complete source code for the ALIGN-2 program is provided in the table below. The ALIGN-2 sequence comparison computer program was authored by Genentech, Inc. and the source code shown in the table below has been filed with user documentation in the U.S. Copyright Office, Washington D.C., 20559, where it is registered under U.S. Copyright Registration No. TXU510087. The ALIGN-2 program is publicly available through Genentech, Inc., South San Francisco, California or may be compiled from the source code provided in the table below. The ALIGN-2 program should be compiled for use on a UNIX operating system, preferably digital UNIX V4.0D. All sequence comparison parameters are set by the ALIGN-2 program and do not vary.

The term "epitope tagged" when used herein refers to a chimeric polypeptide comprising a polypeptide fused to a "tag polypeptide". The tag polypeptide has enough residues to provide an epitope against which an antibody can be made. The tag polypeptide preferably also is fairly unique so that the antibody does not substantially cross-react with other epitopes. Suitable tag polypeptides generally have at least six amino acid residues and usually between about 8 and 50 amino acid residues (preferably, between about 10 and 20 amino acid residues).

As used herein, the term "immunoadhesin" designates antibody-like molecules which combine the binding specificity of a heterologous protein (an "adhesin") with the effector functions of immunoglobulin constant domains. Structurally, the immunoadhesins comprise a fusion of an amino acid sequence with the desired binding specificity which is other than the antigen recognition and binding site of an antibody (i.e., is "heterologous"), and an immunoglobulin constant domain sequence. The adhesin part of an immunoadhesin molecule typically is a contiguous amino acid sequence comprising at least the binding site of a receptor or a ligand. The immunoglobulin constant domain sequence in the immunoadhesin may be obtained from any immunoglobulin, such as IgG-1, IgG-2, IgG-3, or IgG-4 subtypes, IgA (including IgA-1 and IgA-2), IgE, IgD or IgM.

The term "antagonist" is used in the broadest sense, and includes any molecule that partially or fully blocks, inhibits, or neutralizes one or more biological activities of WISP-1 polypeptide, *in vitro*, *in situ*, or *in*

in vivo. Examples of such biological activities of WISP-1 polypeptides include induction or secretion of HAS2, HA, CD44 or RHAMM in at least one type of mammalian cell. An antagonist may function in a direct or indirect manner. For instance, the antagonist may function to partially or fully block, inhibit or neutralize one or more biological activities of WISP-1 polypeptide, *in vitro*, *in situ*, or *in vivo*, e.g., as a result of its direct binding to WISP-1 polypeptide. The antagonist may also function indirectly to partially or fully block, inhibit or neutralize one or more biological activities of WISP-1 polypeptide, *in vitro*, *in situ*, or *in vivo* as a result of, e.g., blocking or inhibiting another effector molecule.

The term "WISP-1 antagonist" refers to any molecule that partially or fully blocks, inhibits, or neutralizes a biological activity of WISP-1 and include but are not limited to, antibodies, immunoadhesins, WISP-1 immunoadhesins, WISP-1 fusion proteins, covalently modified forms of WISP-1, WISP-1 variants and fusion proteins thereof, WISP-1 antibodies, and higher oligomer forms of WISP-1 (dimers, aggregates) or homo- or heteropolymer forms of WISP-1. To determine whether a WISP-1 antagonist molecule partially or fully blocks, inhibits or neutralizes a biological activity of WISP-1, assays may be conducted to assess the effect(s) of the antagonist molecule on, for example, various cells (as described in the Examples) or in an *in vivo* murine model of lung cancer metastasis. The various assays may be conducted in known *in vitro* or *in vivo* assay formats. Preferably, the WISP-1 antagonists employed in the methods described herein will be capable of blocking or neutralizing at least one type of WISP-1 activity, which may optionally be determined in assays such as described herein.

The term "antibody" is used in the broadest sense and specifically covers, for example, single monoclonal antibodies, antibody compositions with polyepitopic specificity, single chain antibodies, and fragments of antibodies. "Antibody" as used herein includes intact immunoglobulin or antibody molecules, polyclonal antibodies, multispecific antibodies (i.e., bispecific antibodies formed from at least two intact antibodies) and immunoglobulin fragments (such as Fab, F(ab')₂, or Fv), so long as they exhibit any of the desired antagonistic properties described herein.

Antibodies are typically proteins or polypeptides which exhibit binding specificity to a specific antigen. Native antibodies are usually heterotetrameric glycoproteins, composed of two identical light (L) chains and two identical heavy (H) chains. Typically, each light chain is linked to a heavy chain by one covalent disulfide bond, while the number of disulfide linkages varies between the heavy chains of different immunoglobulin isotypes. Each heavy and light chain also has regularly spaced intrachain disulfide bridges. Each heavy chain has at one end a variable domain (V_H) followed by a number of constant domains. Each light chain has a variable

domain at one end (V_L) and a constant domain at its other end; the constant domain of the light chain is aligned with the first constant domain of the heavy chain, and the light chain variable domain is aligned with the variable domain of the heavy chain. Particular amino acid residues are believed to form an interface between the light and heavy chain variable domains [Chothia et al., J. Mol. Biol., 186:651-663 (1985); Novotny and Haber, Proc. Natl. Acad. Sci. USA, 82:4592-4596 (1985)]. The light chains of antibodies from any vertebrate species can be assigned to one of two clearly distinct types, called kappa and lambda, based on the amino acid sequences of their constant domains. Depending on the amino acid sequence of the constant domain of their heavy chains, immunoglobulins can be assigned to different classes. There are five major classes of immunoglobulins: IgA, IgD, IgE, IgG and IgM, and several of these may be further divided into subclasses (isotypes), e.g., IgG-1, IgG-2, IgG-3, and IgG-4; IgA-1 and IgA-2. The heavy chain constant domains that correspond to the different classes of immunoglobulins are called alpha, delta, epsilon, gamma, and mu, respectively.

"Antibody fragments" comprise a portion of an intact antibody, generally the antigen binding or variable region of the intact antibody. Examples of antibody fragments include Fab, Fab', F(ab')₂, and Fv fragments, diabodies, single chain antibody molecules, and multispecific antibodies formed from antibody fragments.

The term "variable" is used herein to describe certain portions of the variable domains which differ in sequence among antibodies and are used in the binding and specificity of each particular antibody for its particular antigen. However, the variability is not usually evenly distributed through the variable domains of antibodies. It is typically concentrated in three segments called complementarity determining regions (CDRs) or hypervariable regions both in the light chain and the heavy chain variable domains. The more highly conserved portions of the variable domains are called the framework (FR). The variable domains of native heavy and light chains each comprise four FR regions, largely adopting a β -sheet configuration, connected by three CDRs, which form loops connecting, and in some cases forming part of, the β -sheet structure. The CDRs in each chain are held together in close proximity by the FR regions and, with the CDRs from the other chain, contribute to the formation of the antigen binding site of antibodies [see Kabat, E.A. et al., Sequences of Proteins of Immunological Interest, National Institutes of Health, Bethesda, MD (1987)]. The constant domains are not involved directly in binding an antibody to an antigen, but exhibit various effector functions, such as participation of the antibody in antibody-dependent cellular toxicity.

The term "monoclonal antibody" as used herein refers to an antibody obtained from a population of substantially homogeneous antibodies, i.e., the individual antibodies comprising the population are identical except for possible naturally-occurring mutations that may be present in minor amounts.

5 Monoclonal antibodies are highly specific, being directed against a single antigenic site. Furthermore, in contrast to conventional (polyclonal) antibody preparations which typically include different antibodies directed against different determinants (epitopes), each monoclonal antibody is directed against a single determinant on the antigen.

10 The monoclonal antibodies herein include chimeric, hybrid and recombinant antibodies produced by splicing a variable (including hypervariable) domain of the antibody of interest with a constant domain (e.g. "humanized" antibodies), or a light chain with a heavy chain, or a chain from one species with a chain from another species, or fusions with
15 heterologous proteins, regardless of species of origin or immunoglobulin class or subclass designation, as well as antibody fragments (e.g., Fab, F(ab')₂, and Fv), so long as they exhibit the desired biological activity or properties. See, e.g. U.S. Pat. No. 4,816,567 and Mage et al., in Monoclonal Antibody Production Techniques and Applications, pp.79-97 (Marcel Dekker,
20 Inc.: New York, 1987).

Thus, the modifier "monoclonal" indicates the character of the antibody as being obtained from a substantially homogeneous population of antibodies, and is not to be construed as requiring production of the antibody by any particular method. For example, the monoclonal antibodies to
25 be used in accordance with the present invention may be made by the hybridoma method first described by Kohler and Milstein, Nature, 256:495 (1975), or may be made by recombinant DNA methods such as described in U.S. Pat. No. 4,816,567. The "monoclonal antibodies" may also be isolated from phage libraries generated using the techniques described in McCafferty et al.,
30 Nature, 348:552-554 (1990), for example.

"Humanized" forms of non-human (e.g. murine) antibodies are specific chimeric immunoglobulins, immunoglobulin chains, or fragments thereof (such as Fv, Fab, Fab', F(ab')₂ or other antigen-binding subsequences of antibodies) which contain minimal sequence derived from non-human
35 immunoglobulin. For the most part, humanized antibodies are human immunoglobulins (recipient antibody) in which residues from a complementary determining region (CDR) of the recipient are replaced by residues from a CDR of a non-human species (donor antibody) such as mouse, rat, or rabbit having the desired specificity, affinity, and capacity. In some instances, Fv
40 framework region (FR) residues of the human immunoglobulin are replaced by corresponding non-human residues. Furthermore, the humanized antibody may comprise residues which are found neither in the recipient antibody nor in

the imported CDR or framework sequences. These modifications are made to further refine and optimize antibody performance. In general, the humanized antibody will comprise substantially all of at least one, and typically two, variable domains, in which all or substantially all of the CDR regions correspond to those of a non-human immunoglobulin and all or substantially all of the FR regions are those of a human immunoglobulin consensus sequence. The humanized antibody optimally also will comprise at least a portion of an immunoglobulin constant region or domain (Fc), typically that of a human immunoglobulin.

10 A "human antibody" is one which possesses an amino acid sequence which corresponds to that of an antibody produced by a human and/or has been made using any of the techniques for making human antibodies known in the art or as disclosed herein. This definition of a human antibody includes antibodies comprising at least one human heavy chain polypeptide or at least one human light chain polypeptide, for example an antibody 15 comprising murine light chain and human heavy chain polypeptides. Human antibodies can be produced using various techniques known in the art. In one embodiment, the human antibody is selected from a phage library, where that phage library expresses human antibodies (Vaughan et al. Nature Biotechnology, 14:309-314 (1996); Sheets et al. PNAS, (USA) 95:6157-6162 (1998)); Hoogenboom and Winter, J. Mol. Biol., 227:381 (1991); Marks et al., J. Mol. Biol., 222:581 (1991)). Human antibodies can also be made by introducing human immunoglobulin loci into transgenic animals, e.g., mice 25 in which the endogenous immunoglobulin genes have been partially or completely inactivated. Upon challenge, human antibody production is observed, which closely resembles that seen in humans in all respects, including gene rearrangement, assembly, and antibody repertoire. This approach is described, for example, in U.S. Patent Nos. 5,545,807; 5,545,806; 5,569,825; 5,625,126; 5,633,425; 5,661,016, and in the following 30 scientific publications: Marks et al., Bio/Technology, 10: 779-783 (1992); Lonberg et al., Nature, 368: 856-859 (1994); Morrison, Nature, 368:812-13 (1994); Fishwild et al., Nature Biotechnology, 14: 845-51 (1996); Neuberger, Nature Biotechnology, 14: 826 (1996); Lonberg and Huszar, Intern. Rev. Immunol., 13:65-93 (1995). Alternatively, the human antibody 35 may be prepared via immortalization of human B lymphocytes producing an antibody directed against a target antigen (such B lymphocytes may be recovered from an individual or may have been immunized *in vitro*). See, e.g., Cole et al., Monoclonal Antibodies and Cancer Therapy, Alan R. Liss, p. 77 (1985); Boerner et al., J. Immunol., 147 (1):86-95 (1991); and US Pat 40 No. 5,750,373.

The term "Fc region" is used to define the C-terminal region of an immunoglobulin heavy chain which may be generated by papain digestion of an

A "functional Fc region" possesses at least one "effector function" of a native sequence Fc region. Exemplary "effector functions" include Clq binding; complement dependent cytotoxicity (CDC); Fc receptor binding; antibody-dependent cell-mediated cytotoxicity (ADCC); phagocytosis; down
 5 regulation of cell surface receptors (e.g. B cell receptor; BCR), etc. Such effector functions generally require the Fc region to be combined with a binding domain (e.g. an antibody variable domain) and can be assessed using various assays known in the art for evaluating such antibody effector functions.

10 A "native sequence Fc region" comprises an amino acid sequence identical to the amino acid sequence of a Fc region found in nature. A "variant Fc region" comprises an amino acid sequence which differs from that of a native sequence Fc region by virtue of at least one amino acid modification. Preferably, the variant Fc region has at least one amino
 15 acid substitution compared to a native sequence Fc region or to the Fc region of a parent polypeptide, e.g. from about one to about ten amino acid substitutions, and preferably from about one to about five amino acid substitutions in a native sequence Fc region or in the Fc region of the parent polypeptide. The variant Fc region herein will preferably possess
 20 at least about 80% sequence identity with a native sequence Fc region and/or with an Fc region of a parent polypeptide, and most preferably at least about 90% sequence identity therewith, more preferably at least about 95% sequence identity therewith.

"Antibody-dependent cell-mediated cytotoxicity" and "ADCC" refer to a
 25 cell-mediated reaction in which nonspecific cytotoxic cells that express Fc receptors (FCRs) (e.g. Natural Killer (NK) cells, neutrophils, and macrophages) recognize bound antibody on a target cell and subsequently cause lysis of the target cell. The primary cells for mediating ADCC, NK cells, express FcγRIII only, whereas monocytes express FcγRI, FcγRII and
 30 FcγRIII. FcR expression on hematopoietic cells is summarized in Table 3 on page 464 of Ravetch and Kinet, Annu. Rev. Immunol., 9:457-92 (1991). To assess ADCC activity of a molecule of interest, an *in vitro* ADCC assay, such as that described in US Patent No., 5,500,362 or 5,821,337 may be performed. Useful effector cells for such assays include peripheral blood
 35 mononuclear cells (PBMC) and Natural Killer (NK) cells. Alternatively, or additionally, ADCC activity of the molecule of interest may be assessed *in vivo*, e.g., in a animal model such as that disclosed in Clynes et al. PNAS (USA), 95:652-656 (1998).

"Human effector cells" are leukocytes which express one or more FcRs
 40 and perform effector functions. Preferably, the cells express at least FcγRIII and perform ADCC effector function. Examples of human leukocytes which mediate ADCC include peripheral blood mononuclear cells (PBMC),

Natural killer (NK) cells, monocytes, cytotoxic T cells and neutrophils; with PBMCs and NK cells being preferred. The effector cells may be isolated from a native source thereof, e.g. from blood or PBMCs as described herein.

The terms "Fc receptor" and "FcR" are used to describe a receptor that binds to the Fc region of an antibody. The preferred FcR is a native sequence human FcR. Moreover, a preferred FcR is one which binds an IgG antibody (a gamma receptor) and includes receptors of the FcγRI, FcγRII, and FcγRIII subclasses, including allelic variants and alternatively spliced forms of these receptors. FcγRII receptors include FcγRIIA (an "activating receptor") and FcγRIIB (an "inhibiting receptor"), which have similar amino acid sequences that differ primarily in the cytoplasmic domains thereof. Activating receptor FcγRIIA contains an immunoreceptor tyrosine-based activation motif (ITAM) in its cytoplasmic domain. Inhibiting receptor FcγRIIB contains an immunoreceptor tyrosine-based inhibition motif (ITIM) in its cytoplasmic domain (reviewed in Daeron, Annu. Rev. Immunol., 15:203-234 (1997)). FcRs are reviewed in Ravetch and Kinet, Annu. Rev. Immunol., 9:457-92 (1991); Capel et al., Immunomethods, 4:25-34 (1994); and de Haas et al., J. Lab. Clin. Med., 126:330-41 (1995). Other FcRs, including those to be identified in the future, are encompassed by the term "FcR" herein. The term also includes the neonatal receptor, FcRn, which is responsible for the transfer of maternal IgGs to the fetus (Guyer et al., J. Immunol., 117:587 (1976); and Kim et al., J. Immunol., 24:249 (1994)).

"Complement dependent cytotoxicity" and "CDC" refer to the lysing of a target in the presence of complement. The complement activation pathway is initiated by the binding of the first component of the complement system (C1q) to a molecule (e.g. an antibody) complexed with a cognate antigen. To assess complement activation, a CDC assay, e.g. as described in Gazzano-Santoro et al., J. Immunol. Methods, 202:163 (1996), may be performed.

An "affinity matured" antibody is one with one or more alterations in one or more CDRs thereof which result an improvement in the affinity of the antibody for antigen, compared to a parent antibody which does not possess those alteration(s). Preferred affinity matured antibodies will have nanomolar or even picomolar affinities for the target antigen. Affinity matured antibodies are produced by procedures known in the art. Marks et al. Bio/Technology, 10:779-783 (1992) describes affinity maturation by VH and VL domain shuffling. Random mutagenesis of CDR and/or framework residues is described by: Barbas et al. Proc Nat. Acad. Sci, USA 91:3809-3813 (1994); Schier et al. Gene, 169:147-155 (1995); Yelton et al. J. Immunol., 155:1994-2004 (1995); Jackson et al., J. Immunol., 154(7):3310-9 (1995); and Hawkins et al., J. Mol. Biol., 226:889-896 (1992).

The term "immunospecific" as used in "immunospecific binding of antibodies" for example, refers to the antigen specific binding interaction that occurs between the antigen-combining site of an antibody and the specific antigen recognized by that antibody.

5 The terms "cancer", "cancerous", "metastasis" and "malignant" refer to or describe the physiological condition in mammals that is typically characterized by unregulated cell growth. Examples of cancer include but are not limited to, carcinoma including adenocarcinoma, lymphoma, blastoma, melanoma, sarcoma, and leukemia. More particular examples of such cancers
10 include squamous cell cancer, small-cell lung cancer, non-small cell lung cancer, gastrointestinal cancer, Hodgkin's and non-Hodgkin's lymphoma, pancreatic cancer, glioblastoma, cervical cancer, ovarian cancer, liver cancer such as hepatic carcinoma and hepatoma, bladder cancer, breast cancer, colon cancer, colorectal cancer, endometrial carcinoma, myeloma
15 (such as multiple myeloma), salivary gland carcinoma, kidney cancer such as renal cell carcinoma and Wilms' tumors, basal cell carcinoma, melanoma, prostate cancer, vulval cancer, thyroid cancer, testicular cancer, esophageal cancer, and various types of head and neck cancer. The preferred cancers for treatment herein include breast cancer, gastric
20 cancer, lung cancer, colon or colorectal cancer, glioma and glioblastoma.

The term "prodrug" as used in this application refers to a precursor or derivative form of a pharmaceutically active substance that is less cytotoxic to cancer cells compared to the parent drug and is capable of being enzymatically activated or converted into the more active parent
25 form. See, e.g., Wilman, "Prodrugs in Cancer Chemotherapy" *Biochemical Society Transactions*, 14, pp. 375-382, 615th Meeting Belfast (1986) and Stella et al., "Prodrugs: A Chemical Approach to Targeted Drug Delivery," *Directed Drug Delivery*, Borchardt et al., (ed.), pp. 247-267, Humana Press (1985). The prodrugs of this invention include, but are not limited to,
30 phosphate-containing prodrugs, thiophosphate-containing prodrugs, sulfate-containing prodrugs, peptide-containing prodrugs, D-amino acid-modified prodrugs, glycosylated prodrugs, beta-lactam-containing prodrugs, optionally substituted phenoxyacetamide-containing prodrugs or optionally substituted phenylacetamide-containing prodrugs, 5-fluorocytosine and other
35 5-fluorouridine prodrugs which can be converted into the more active cytotoxic free drug. Examples of cytotoxic drugs that can be derivatized into a prodrug form for use in this invention include, but are not limited to, those chemotherapeutic agents described below.

The term "cytotoxic agent" as used herein refers to a substance that
40 inhibits or prevents the function of cells and/or causes destruction of cells. The term is intended to include radioactive isotopes (e.g. At²¹¹, I¹³¹, I¹²⁵, Y⁹⁰, Re¹⁸⁶, Re¹⁸⁸, Sm¹⁵³, Bi²¹², P³² and radioactive isotopes of Lu),

chemotherapeutic agents, and toxins such as small molecule toxins or enzymatically active toxins of bacterial, fungal, plant or animal origin, including fragments and/or variants thereof.

A "chemotherapeutic agent" is a chemical compound useful in the treatment of conditions like cancer. Examples of chemotherapeutic agents include alkylating agents such as thiotepa and cyclophosphamide (CYTOXANTM); alkyl sulfonates such as busulfan, improsulfan and piposulfan; aziridines such as benzodopa, carboquone, meturedopa, and uredopa; ethylenimines and methylamelamines including altretamine, triethylenemelamine, triethylenephosphoramide, triethylenethiophosphoramide and trimethylolomelamine; acetogenins (especially bullatacin and bullatacinone); a camptothecin (including the synthetic analogue topotecan); bryostatin; callystatin; CC-1065 (including its adozelesin, carzelesin and bizelesin synthetic analogues); cryptophycins (particularly cryptophycin 1 and cryptophycin 8); dolastatin; duocarmycin (including the synthetic analogues, KW-2189 and CBI-TMI); eleutherobin; pancratistatin; a sarcodictyin; spongistatin; nitrogen mustards such as chlorambucil, chlornaphazine, cholophosphamide, estramustine, ifosfamide, mechlorethamine, mechlorethamine oxide hydrochloride, melphalan, novembichin, phenesterine, prednimustine, trofosfamide, uracil mustard; nitrosureas such as carmustine, chlorozotocin, fotemustine, lomustine, nimustine, ranimustine; antibiotics such as the enediyne antibiotics (e.g. calicheamicin, especially calicheamicin γ_1^I and calicheamicin θ_1^I , see, e.g., Agnew Chem Intl. Ed. Engl., 33:183-186 (1994); dynemicin, including dynemicin A; an esperamicin; as well as neocarzinostatin chromophore and related chromoprotein enediyne antibiotic chromophores), aclacinomysins, actinomycin, authramycin, azaserine, bleomycins, cactinomycin, carabycin, carminomycin, carzinophilin, chromomycins, dactinomycin, daunorubicin, detorubicin, 6-diazo-5-oxo-L-norleucine, doxorubicin (including morpholino-doxorubicin, cyanomorpholino-doxorubicin, 2'-pyrrolino-doxorubicin and deoxydoxorubicin), epirubicin, esorubicin, idarubicin, marcellomycin, mitomycins, mycophenolic acid, nogalamycin, olivomycins, peplomycin, potfiromycin, puromycin, quelamycin, rodorubicin, streptonigrin, streptozocin, tubercidin, ubenimex, zinostatin, zorubicin; anti-metabolites such as methotrexate and 5-fluorouracil (5-FU); folic acid analogues such as denopterin, methotrexate, pteropterin, trimetrexate; purine analogs such as fludarabine, 6-mercaptopurine, thiamiprine, thioguanine; pyrimidine analogs such as ancitabine, azacitidine, 6-azauridine, carmofur, cytarabine, dideoxyuridine, doxifluridine, enocitabine, floxuridine, 5-FU; androgens such as calusterone, dromostanolone propionate, epitioestanol, mepitioestane, testolactone; anti-adrenals such as aminoglutethimide, mitotane, trilostane; folic acid replenisher such as frolinic acid;

aceglatone; aldophosphamide glycoside; aminolevulinic acid; amsacrine;
 bestrabucil; bisantrene; edatraxate; defofamine; demecolcine; diaziquone;
 elfornithine; elliptinium acetate; an epothilone; etoglucid; gallium
 nitrate; hydroxyurea; lentinan; lonidamine; maytansinoids such as
 5 maytansine and ansamitocins; mitoguazone; mitoxantrone; mopidamol;
 nitracrine; pentostatin; phenamet; pirarubicin; podophyllinic acid; 2-
 ethylhydrazide; procarbazine; PSK®; razoxane; rhizoxin; sizofiran;
 spirogermanium; tenuazonic acid; triaziquone; 2, 2', 2''-
 trichlorotriethylamine; trichothecenes (especially T-2 toxin, verracurin A,
 10 roridin A and anguidine); urethan; vindesine; dacarbazine; mannomustine;
 mitobronitol; mitolactol; pipobroman; gacytosine; arabinoside ("Ara-C");
 cyclophosphamide; thiotepa; taxoids, e.g. paclitaxel (TAXOL®, Bristol-Myers
 Squibb Oncology, Princeton, NJ) and doxetaxel (TAXOTERE®, Rhône-Poulenc
 Rorer, Antony, France); chlorambucil; gemcitabine; 6-thioguanine;
 15 mercaptopurine; methotrexate; platinum analogs such as cisplatin and
 carboplatin; vinblastine; platinum; etoposide (VP-16); ifosfamide;
 mitomycin C; mitoxantrone; vincristine; vinorelbine; navelbine; novantrone;
 teniposide; daunomycin; aminopterin; xeloda; ibandronate; CPT-11;
 topoisomerase inhibitor RFS 2000; difluoromethylornithine (DMFO); retinoic
 20 acid; capecitabine; and pharmaceutically acceptable salts, acids or
 derivatives of any of the above. Also included in this definition are
 anti-hormonal agents that act to regulate or inhibit hormone action on
 tumors such as anti-estrogens including for example tamoxifen, raloxifene,
 aromatase inhibiting 4(5)-imidazoles, 4-hydroxytamoxifen, trioxifene,
 25 keoxifene, LY117018, onapristone, and toremifene (Fareston); and anti-
 androgens such as flutamide, nilutamide, bicalutamide, leuprolide, and
 goserelin; and pharmaceutically acceptable salts, acids or derivatives of
 any of the above.

A "growth inhibitory agent" when used herein refers to a compound or
 30 composition which inhibits growth of a cell, either *in vitro* or *in vivo*.
 Thus, the growth inhibitory agent is one which significantly reduces the
 percentage of cells overexpressing such genes in S phase. Examples of
 growth inhibitory agents include agents that block cell cycle progression
 (at a place other than S phase), such as agents that induce G1 arrest and
 35 M-phase arrest. Classical M-phase blockers include the vincas (vincristine
 and vinblastine), taxol, and topo II inhibitors such as doxorubicin,
 epirubicin, daunorubicin, etoposide, and bleomycin. Those agents that
 arrest G1 also spill over into S-phase arrest, for example, DNA alkylating
 agents such as tamoxifen, prednisone, dacarbazine, mechlorethamine,
 40 cisplatin, methotrexate, 5-fluorouracil, and ara-C. Further information
 can be found in *The Molecular Basis of Cancer*, Mendelsohn and Israel, eds.,
 Chapter 1, entitled "Cell cycle regulation, oncogens, and antineoplastic

drugs" by Murakami et al. (WB Saunders: Philadelphia, 1995), especially p. 13.

The term "cytokine" is a generic term for proteins released by one cell population which act on another cell as intercellular mediators. Examples of such cytokines are lymphokines, monokines, and traditional polypeptide hormones. Included among the cytokines are growth hormone such as human growth hormone, N-methionyl human growth hormone, and bovine growth hormone; parathyroid hormone; thyroxine; insulin; proinsulin; relaxin; prorelaxin; glycoprotein hormones such as follicle stimulating hormone (FSH), thyroid stimulating hormone (TSH), and luteinizing hormone (LH); hepatic growth factor; fibroblast growth factor; prolactin; placental lactogen; tumor necrosis factor- α and - β ; Apo-2 ligand (also referred to as TRAIL); mullerian-inhibiting substance; mouse gonadotropin-associated peptide; inhibin; activin; vascular endothelial growth factor; integrin; thrombopoietin (TPO); nerve growth factors; platelet-growth factor; transforming growth factors (TGFs) such as TGF- α and TGF- β ; insulin-like growth factor-I and -II; erythropoietin (EPO); osteoinductive factors; interferons such as interferon- α , - β , and - γ ; colony stimulating factors (CSFs) such as macrophage-CSF (M-CSF); granulocyte-macrophage-CSF (GM-CSF); and granulocyte-CSF (G-CSF); interleukins (ILs) such as IL-1, IL-2, IL-3, IL-4, IL-5, IL-6, IL-7, IL-8, IL-9, IL-11, IL-12; and other polypeptide factors including LIF and kit ligand (KL). As used herein, the term cytokine includes proteins from natural sources or from recombinant cell culture and biologically active equivalents of the native sequence cytokines.

"Treatment" is an intervention performed with the intention of preventing the development or altering the pathology of a disorder. Accordingly, "treatment" refers to both therapeutic treatment and prophylactic or preventative measures, wherein the object is to prevent or slow down (lessen) the targeted pathological condition or disorder. Those in need of treatment include those already with the disorder as well as those in which the disorder is to be prevented. In treatment of a disorder, a therapeutic agent may directly decrease or increase the magnitude of response of a pathological component of the disorder, or render the disease more susceptible to treatment by other therapeutic agents, e.g. antibiotics, antifungals, anti-inflammatory agents, chemotherapeutics, etc.

The term "effective amount" is the minimum concentration of WISP-1 antagonist which causes, induces or results in either a detectable improvement or reduction of a pathological condition. In a method of treating cancer, an effective amount is one which causes, induces, or results in reduction of cancer cell number or tumor volume. Furthermore a

"therapeutically effective amount" is the minimum concentration (amount) of WISP-1 antagonist administered to a mammal which would be effective in at least attenuating a pathological symptom.

"Chronic" administration refers to administration of the factor(s) in a continuous mode as opposed to an acute mode, so as to maintain the initial therapeutic effect (activity) for an extended period of time. "Intermittent" administration is treatment that is done not consecutively without interruption, but rather is cyclic in nature.

10 "Mammal" for purposes of treatment refers to any animal classified as a mammal, including humans, domestic and farm animals, and zoo, sports, or pet animals, such as dogs, horses, cats, cattle, pigs, hamsters, etc. Preferably, the mammal is human.

Administration "in combination with" one or more further therapeutic agents includes simultaneous (concurrent) and consecutive administration in any order.

any order.

"Carriers" as used herein include pharmaceutically acceptable carriers, excipients, or stabilizers which are nontoxic to the cell or mammal being exposed thereto at the dosages and concentrations employed. Often the physiologically acceptable carrier is an aqueous pH buffered solution. Examples of physiologically acceptable carriers include buffers such as phosphate, citrate, and other organic acids; antioxidants including ascorbic acid; low molecular weight (less than about 10 residues) polypeptide; proteins, such as serum albumin, gelatin, or immunoglobulins; hydrophilic polymers such as polyvinylpyrrolidone; amino acids such as glycine, glutamine, asparagine, arginine or lysine; monosaccharides, disaccharides, and other carbohydrates including glucose, mannose, or dextrans; chelating agents such as EDTA; sugar alcohols such as mannitol or sorbitol; salt-forming counterions such as sodium; and/or nonionic surfactants such as TWEEN®, polyethylene glycol (PEG), and PLURONICS®, hyaluronic acid (HA).

II. Methods and Compositions of the Invention

As described further in the Examples below, Applicant has surprisingly found that ectopic expression of WISP-1 in fibroblasts triggered HA production, namely increased HA production induced its accumulation at the cell surface and in the culture media. It is believed that cell surface HA can enhance anchorage-independent growth and tumorigenicity (Kosaki et al., Cancer Res., 59:1141-1145 (1999)) and may diminish contact inhibition of cell proliferation, thereby promoting cell migration (Ichikawa et al., J. Invest. Dermatol., 113:935-939 (1999); Itano et al., Proc. Natl. Acad. Sci., 99:3609-3614 (2002)). Hyaluronan coat formation has also been correlated with cellular metastatic potential

Zhang et al., Cancer Res., 55:428-433 (1995); Toole et al., J. Biol. Chem., 277:4593-4596 (2002)). Analysis of HA synthases (HAS1, HAS2, and HAS3) expression revealed that WISP-1 promoted HAS2 induction whereas HAS1 and HAS3 mRNA levels remained unchanged. HAS2 is the main target of growth factor and cytokine up-regulated HA synthesis (Pienimäki et al., J. Biol. Chem., 276:20428-20435 (2001)). Ectopic expression of HAS2 increases HA secretion and induces hyaluronan coat formation (Kosaki et al., supra). Moreover, HAS2 is believed to be important for hyaluronan-mediated transformation of epithelium to mesenchyme during cardiac morphogenesis (Camenisch et al., J. Clin. Invest., 106:349-360 (2000)). Because epithelial to mesenchymal transition is also important for tumor invasion and metastasis, HAS2 is postulated to play an important role in epithelial tumor progression (Boyer et al., Biochem. Pharmacol., 60:1091-1099 (2000); Hay, Acta Anat., 154:8-20 (1995); Arias, Cell, 105:425-431 (2001)). These results indicate that HA secretion induced by WISP-1 may be important for tumor invasion and metastasis.

WISP-1 also induced the expression of two HA receptors, CD44 and RHAMM. By inducing HA receptor expression and increasing HA production, WISP-1 may activate an autocrine and/or paracrine loop. Hyaluronan interaction with CD44 and RHAMM promotes cell locomotion and proliferation *in vitro* and tumor growth and metastasis *in vivo* (Turley et al., J. Biol. Chem., 277:4589-4592 (2002); Sy et al., Curr. Top. Microbiol. Immunol., 213:129-153 (1996); Hall et al., J. Neurooncol., 26:221-229 (1995)).

The effects of WISP-1 on cell migration was analyzed in a cell wound healing assay and by time lapse microscopy, and the data showed that WISP-1 expression promoted cell motility. Further, isolated cells showed increased migration suggesting that WISP-1 could act through an autocrine mechanism. Purified recombinant WISP-1 promoted HAS2 and CD44 expression, though, when attached to a surface. This induction was further increased when WISP-1 was tethered through its interaction with decorin. Similarly, WISP-1 promoted cell motility when surface bound, suggesting that WISP-1 may also act through a paracrine mechanism when tethered to a substrate. The Examples herein show that WISP-1 induced cell motility was mediated by CD44, by abolishing migration with an anti-CD44 antibody. WISP-1 induced cell motility was also inhibited by WISP-1 antibodies. Moreover, WISP-1 haptotactic activity was not restricted to a single cell type as it induced both normal fibroblast and colon adenocarcinoma cell migration.

Because WISP-1 is believed to be a Wnt-1 downstream effector, the C57MG/Wnt-1 cell line was analyzed for phenotypes found in NRK/WISP-1 cells. Consistent with a role for WISP-1 downstream of Wnt-1, C57MG/Wnt-1 cells overexpressed HAS2 and CD44 and had higher CD44 protein content compared to the parental cell line. In addition, C57/Wnt-1 cells

spontaneously scattered in culture and demonstrated a de-differentiated spindloid morphology similar to NRK/WISP-1H cells.

Expression analysis was also performed on a group of MMTV-Wnt-1 mammary tumors, and elevated CD44 and HAS2 expression was detected in all mammary tumors from the MMTV-Wnt-1 transgenic mice. In these tumors, WISP-1 expression was localized to the stromal fibroblasts whereas CD44 and HAS2 were expressed by tumor epithelial cells. Although negative for HAS2 expression, the peritumoral stroma contained high levels of HA whereas the tumor parenchyma stained only weakly for HA. Although it is generally accepted that fibroblasts are responsible for HA production, the origin of stromal HA is not definitely known. Because HAS2 expression was found only in the parenchyma, the experimental results disclosed herein suggest that the tumor cells are responsible for the HA synthesis and deposition in the stroma. Hyaluronan accumulation in the peritumoral stroma is frequently encountered in several tumor types and was previously reported for ovary, breast, prostate and colon adenocarcinomas (Ropponen et al., Cancer Res., 58:342-347 (1998); Lipponen et al., Eur. J. Cancer, 37:849-856 (2001); Auvinen et al., Am. J. Pathol., 156:529-536 (2000); Anttila et al., Cancer Res., 60:150-155 (2000)). Moreover, a high stromal level of HA was associated with poor differentiation, metastatic behavior and unfavorable prognosis.

Because HA and CD44 are associated with tumor invasion, the metastatic potential of WISP-1 expressing cells was evaluated in vivo. After tail inoculation, NRK/WISP-1 cells readily colonized the lungs of injected mice and formed invasive tumors. NRK/WISP-1 cells exhibited a significant metastatic potential while NRK cells did not. Histological observation revealed that NRK/WISP-1 cells populated the vasculature and invaded the pulmonary airways. The lung colonization was proportional to the number of cells injected, the time after injection and the WISP-1 expression. In addition, Applicant found the metastatic potential to be proportional to CD44 and HAS2 expression levels, and the treatment of mice inoculated with NRK/WISP-1H cells with CD44 antibody or WISP-1 antibody greatly diminished the number and size of the tumors in the lung. Thus, it is likely that levels of WISP-1 promoted lung colonization through a hyaluronan-CD44 mechanism. This is consistent with previous reports demonstrating the importance of CD44 and its interaction with HA for tumor growth, metastasis and the retention of metastatic cells to the lung vasculature (Sy et al., supra; Kogerman et al., Proc. Natl. Acad. Sci., 94:13233-13238 (1997)).

Although activation of the Wnt pathway by APC or β -catenin mutations may typically be associated with colorectal cancer, several lines of evidence suggest that it also plays a role in other types of cancer

including mammary adenocarcinoma (Polakis, Genes Dev., 14:1837-1851 (2000); Brown, Breast Cancer Res., 3:351-355 (2001)). APC truncation and increased β -catenin levels were found in human breast cancer cell lines (Schlosshauer et al., Carcinogenesis, 21:1453-1456 (2000)). Somatic mutations of the APC gene were found in primary breast cancers (Furuuchi et al., Am. J. Pathol., 156:1997-2005 (2000)). In addition, β -catenin activating mutations promote mouse mammary adenocarcinomas Michaelson et al., Oncogene, 20:5525-5532 (2001)). Elevated levels of Wnt-1 and β -catenin were found in invasive ductal breast carcinomas and correlated with poor prognosis (Lin et al., Proc. Natl. Acad. Sci., 97:4262-4266 (2000)). It is therefore possible that WISP-1 expression found in certain breast adenocarcinomas resulted from Wnt pathway activation. Applicants have also found, by Taqman analysis, that WISP-1 is overexpressed in cancers such as breast and glial tumors.

For at least these reasons, it is believed that WISP-1 antagonists will be particularly useful in treating and diagnosing various pathological disorders, such as cancer. The present invention accordingly provides methods for modulating, blocking or neutralizing WISP-1 activity in mammalian cells which comprise exposing the cells to a desired amount of WISP-1 antagonist. Preferably, the amount of WISP-1 antagonist employed will be an amount effective to reduce or inhibit cancer cell growth, metastasis or motility. This can be accomplished in vivo or ex vivo in accordance, for instance, with the methods described below and in the Examples. Exemplary conditions or disorders to be treated with such WISP-1 antagonists include conditions in mammals clinically referred to as cancer.

Diagnostic methods are also provided herein. For instance, the antagonists may be employed to detect invasive or metastatic cancers. The antagonist molecule may be used, e.g., in assays to detect or quantitate metastatic cancer cells in a sample. A sample, such as cells obtained from a mammal, can be incubated in the presence of a labeled antagonist, and detection of the labeled antagonist bound in the sample can be performed.

The antagonists which can be employed in the methods include, but are not limited to, WISP-1 immunoadhesins, fusion proteins comprising WISP-1, covalently modified forms of WISP-1, WISP-1 variants, fusion proteins thereof, and WISP-1 antibodies. Various techniques that can be employed for making the antagonists are described herein. For instance, methods and techniques for preparing WISP-1 polypeptides are described. Further modifications of the polypeptides, and antibodies to WISP-1 are also described.

In addition to the full-length native sequence WISP-1 polypeptide described herein, it is contemplated that WISP-1 polypeptide variants can be prepared. WISP-1 variants can be prepared by introducing appropriate

nucleotide changes into the encoding DNA, and/or by synthesis of the desired polypeptide. Those skilled in the art will appreciate that amino acid changes may alter post-translational processes of the WISP-1 polypeptide, such as changing the number or position of glycosylation sites or altering the membrane anchoring characteristics.

Variations in the WISP-1 polypeptides described herein, can be made, for example, using any of the techniques and guidelines for conservative and non-conservative mutations set forth, for instance, in U.S. Patent No. 5,364,934. Variations may be a substitution, deletion or insertion of one or more codons encoding the polypeptide that results in a change in the amino acid sequence as compared with the native sequence polypeptide. Optionally the variation is by substitution of at least one amino acid with any other amino acid in one or more of the domains of the WISP-1 polypeptide. Guidance in determining which amino acid residue may be inserted, substituted or deleted without adversely affecting the desired activity may be found by comparing the sequence of the WISP-1 polypeptide with that of homologous known protein molecules and minimizing the number of amino acid sequence changes made in regions of high homology. Amino acid substitutions can be the result of replacing one amino acid with another amino acid having similar structural and/or chemical properties, such as the replacement of a leucine with a serine, i.e., conservative amino acid replacements. Insertions or deletions may optionally be in the range of about 1 to 5 amino acids. The variation allowed may be determined by systematically making insertions, deletions or substitutions of amino acids in the sequence and testing the resulting variants for activity exhibited by the full-length or mature native sequence.

WISP-1 polypeptide fragments are provided herein. Such fragments may be truncated at the N-terminus or C-terminus, or may lack internal residues, for example, when compared with a full length native protein. Certain fragments lack amino acid residues that are not essential for a desired biological activity of the WISP-1 polypeptide.

WISP-1 polypeptide fragments may be prepared by any of a number of conventional techniques. Desired peptide fragments may be chemically synthesized. An alternative approach involves generating polypeptide fragments by enzymatic digestion, e.g., by treating the protein with an enzyme known to cleave proteins at sites defined by particular amino acid residues, or by digesting the DNA with suitable restriction enzymes and isolating the desired fragment. Yet another suitable technique involves isolating and amplifying a DNA fragment encoding a desired polypeptide fragment, by polymerase chain reaction (PCR). Oligonucleotides that define the desired termini of the DNA fragment are employed at the 5' and 3' primers in the PCR.

In particular embodiments, conservative substitutions of interest are shown in the Table below under the heading of preferred substitutions. If such substitutions result in a change in biological activity, then more substantial changes, denominated exemplary substitutions in the Table, or as further described below in reference to amino acid classes, are introduced and the products screened.

<u>Table</u>			
	<u>Original</u>	<u>Exemplary</u>	<u>Preferred</u>
	<u>Residue</u>	<u>Substitutions</u>	<u>Substitutions</u>
10	Ala (A)	val; leu; ile	val
	Arg (R)	lys; gln; asn	lys
	Asn (N)	gln; his; lys; arg	gln
	Asp (D)	glu	glu
	Cys (C)	ser	ser
15	Gln (Q)	asn	asn
	Glu (E)	asp	asp
	Gly (G)	pro; ala	ala
	His (H)	asn; gln; lys; arg	arg
20	Ile (I)	leu; val; met; ala; phe; norleucine	leu
	Leu (L)	norleucine; ile; val; met; ala; phe	ile
	Lys (K)	arg; gln; asn	arg
25	Met (M)	leu; phe; ile	leu
	Phe (F)	leu; val; ile; ala; tyr	leu
	Pro (P)	ala	ala
	Ser (S)	thr	thr
	Thr (T)	ser	ser
30	Trp (W)	tyr; phe	tyr
	Tyr (Y)	trp; phe; thr; ser	phe
	Val (V)	ile; leu; met; phe; ala; norleucine	leu

Substantial modifications in function or immunological identity of the WISP-1 polypeptide are accomplished by selecting substitutions that differ significantly in their effect on maintaining (a) the structure of the polypeptide backbone in the area of the substitution, for example, as a sheet or helical conformation, (b) the charge or hydrophobicity of the molecule at the target site, or (c) the bulk of the side chain. Naturally occurring residues are divided into groups based on common side-chain properties:

- (1) hydrophobic: norleucine, met, ala, val, leu, ile;
- (2) neutral hydrophilic: cys, ser, thr;
- (3) acidic: asp, glu;
- (4) basic: asn, gln, his, lys, arg;
- 5 (5) residues that influence chain orientation: gly, pro; and
- (6) aromatic: trp, tyr, phe.

Non-conservative substitutions will entail exchanging a member of one of these classes for another class. Such substituted residues also may be introduced into the conservative substitution sites or, more preferably, 10 into the remaining (non-conserved) sites.

The variations can be made using methods known in the art such as oligonucleotide-mediated (site-directed) mutagenesis, alanine scanning, and PCR mutagenesis. Site-directed mutagenesis [Carter et al., Nucl. Acids Res., 13:4331 (1986); Zoller et al., Nucl. Acids Res., 10:6487 (1987)], 15 cassette mutagenesis [Wells et al., Gene, 34:315 (1985)], restriction selection mutagenesis [Wells et al., Philos. Trans. R. Soc. London SerA, 317:415 (1986)] or other known techniques can be performed on the cloned DNA to produce the WISP-1 polypeptide variant DNA.

Scanning amino acid analysis can also be employed to identify one or 20 more amino acids along a contiguous sequence. Among the preferred scanning amino acids are relatively small, neutral amino acids. Such amino acids include alanine, glycine, serine, and cysteine. Alanine is typically a preferred scanning amino acid among this group because it eliminates the side-chain beyond the beta-carbon and is less likely to alter the main- 25 chain conformation of the variant [Cunningham and Wells, Science, 244:1081-1085 (1989)]. Alanine is also typically preferred because it is the most common amino acid. Further, it is frequently found in both buried and exposed positions [Creighton, The Proteins, (W.H. Freeman & Co., N.Y.); Chothia, J. Mol. Biol., 150:1 (1976)]. If alanine substitution does not 30 yield adequate amounts of variant, an isoteric amino acid can be used.

Any cysteine residue not involved in maintaining the proper conformation of the WISP-1 polypeptide also may be substituted, generally with serine, to improve the oxidative stability of the molecule and prevent aberrant crosslinking. Conversely, cysteine bond(s) may be added to the 35 WISP-1 polypeptide to improve its stability.

The description below relates primarily to production of WISP-1 polypeptides by culturing cells transformed or transfected with a vector containing WISP-1 polypeptide-encoding nucleic acid. It is, of course, contemplated that alternative methods, which are well known in the art, may 40 be employed to prepare WISP-1 polypeptides. For instance, the appropriate amino acid sequence, or portions thereof, may be produced by direct peptide synthesis using solid-phase techniques [see, e.g., Stewart et al., Solid-

Phase Peptide Synthesis, W.H. Freeman Co., San Francisco, CA (1969); Merrifield, J. Am. Chem. Soc., 85:2149-2154 (1963)]. In vitro protein synthesis may be performed using manual techniques or by automation. Automated synthesis may be accomplished, for instance, using an Applied Biosystems Peptide Synthesizer (Foster City, CA) using manufacturer's instructions. Various portions of the WISP-1 polypeptide may be chemically synthesized separately and combined using chemical or enzymatic methods to produce the desired WISP-1 polypeptide. The methods and techniques described are similarly applicable to production of WISP-1 variants, modified forms of WISP-1 and WISP-1 antibodies.

1. Isolation of DNA Encoding WISP-1 Polypeptide

DNA encoding WISP-1 polypeptide may be obtained from a cDNA library prepared from tissue believed to possess the WISP-1 polypeptide mRNA and to express it at a detectable level. Accordingly, human WISP-1 polypeptide DNA can be conveniently obtained from a cDNA library prepared from human tissue. The WISP-1 polypeptide-encoding gene may also be obtained from a genomic library or by known synthetic procedures (e.g., automated nucleic acid synthesis).

Libraries can be screened with probes (such as oligonucleotides of at least about 20-80 bases) designed to identify the gene of interest or the protein encoded by it. Screening the cDNA or genomic library with the selected probe may be conducted using standard procedures, such as described in Sambrook et al., Molecular Cloning: A Laboratory Manual (New York: Cold Spring Harbor Laboratory Press, 1989). An alternative means to isolate the gene encoding WISP-1 polypeptide is to use PCR methodology [Sambrook et al., supra; Dieffenbach et al., PCR Primer: A Laboratory Manual (Cold Spring Harbor Laboratory Press, 1995)].

Techniques for screening a cDNA library are well known in the art. The oligonucleotide sequences selected as probes should be of sufficient length and sufficiently unambiguous that false positives are minimized. The oligonucleotide is preferably labeled such that it can be detected upon hybridization to DNA in the library being screened. Methods of labeling are well known in the art, and include the use of radiolabels like ³²P-labeled ATP, biotinylation or enzyme labeling. Hybridization conditions, including moderate stringency and high stringency, are provided in Sambrook et al., supra.

Sequences identified in such library screening methods can be compared and aligned to other known sequences deposited and available in public databases such as GenBank or other private sequence databases. Sequence identity (at either the amino acid or nucleotide level) within defined regions of the molecule or across the full-length sequence can be determined using methods known in the art and as described herein.

Nucleic acid having protein coding sequence may be obtained by screening selected cDNA or genomic libraries using the deduced amino acid sequence disclosed herein for the first time, and, if necessary, using conventional primer extension procedures as described in Sambrook et al., supra, to detect precursors and processing intermediates of mRNA that may not have been reverse-transcribed into cDNA.

2. Selection and Transformation of Host Cells

Host cells are transfected or transformed with expression or cloning vectors described herein for WISP-1 polypeptide production and cultured in conventional nutrient media modified as appropriate for inducing promoters, selecting transformants, or amplifying the genes encoding the desired sequences. The culture conditions, such as media, temperature, pH and the like, can be selected by the skilled artisan without undue experimentation.

In general, principles, protocols, and practical techniques for maximizing the productivity of cell cultures can be found in Mammalian Cell Biotechnology: a Practical Approach, M. Butler, ed. (IRL Press, 1991) and Sambrook et al., supra.

Methods of eukaryotic cell transfection and prokaryotic cell transformation are known to the ordinarily skilled artisan, for example, CaCl_2 , CaPO_4 , liposome-mediated and electroporation. Depending on the host cell used, transformation is performed using standard techniques appropriate to such cells. The calcium treatment employing calcium chloride, as described in Sambrook et al., supra, or electroporation is generally used for prokaryotes. Infection with *Agrobacterium tumefaciens* is used for transformation of certain plant cells, as described by Shaw et al., Gene, 23:315 (1983) and WO 89/05859 published 29 June 1989. For mammalian cells without such cell walls, the calcium phosphate precipitation method of Graham and van der Eb, Virology, 52:456-457 (1978) can be employed. General aspects of mammalian cell host system transfections have been described in U.S. Patent No. 4,399,216. Transformations into yeast are typically carried out according to the method of Van Solingen et al., J. Bact., 130:946 (1977) and Hsiao et al., Proc. Natl. Acad. Sci. (USA), 76:3829 (1979). However, other methods for introducing DNA into cells, such as by nuclear microinjection, electroporation, bacterial protoplast fusion with intact cells, or polycations, e.g., polybrene, polyornithine, may also be used. For various techniques for transforming mammalian cells, see Keown et al., Methods in Enzymology, 185:527-537 (1990) and Mansour et al., Nature, 336:348-352 (1988).

Suitable host cells for cloning or expressing the DNA in the vectors herein include prokaryote, yeast, or higher eukaryote cells. Suitable prokaryotes include but are not limited to eubacteria, such as Gram-

negative or Gram-positive organisms, for example, Enterobacteriaceae such as *E. coli*. Various *E. coli* strains are publicly available, such as *E. coli* K12 strain MM294 (ATCC 31,446); *E. coli* X1776 (ATCC 31,537); *E. coli* strain W3110 (ATCC 27,325) and K5 772 (ATCC 53,635). Other suitable prokaryotic host cells include Enterobacteriaceae such as *Escherichia*, e.g., *E. coli*, *Enterobacter*, *Erwinia*, *Klebsiella*, *Proteus*, *Salmonella*, e.g., *Salmonella typhimurium*, *Serratia*, e.g., *Serratia marcescens*, and *Shigella*, as well as *Bacilli* such as *B. subtilis* and *B. licheniformis* (e.g., *B. licheniformis* 41P disclosed in DD 266,710 published 12 April 1989), *Pseudomonas* such as *P. aeruginosa*, and *Streptomyces*. These examples are illustrative rather than limiting. Strain W3110 is one particularly preferred host or parent host because it is a common host strain for recombinant DNA product fermentations. Preferably, the host cell secretes minimal amounts of proteolytic enzymes. For example, strain W3110 may be modified to effect a genetic mutation in the genes encoding proteins endogenous to the host, with examples of such hosts including *E. coli* W3110 strain 1A2, which has the complete genotype *tonA*; *E. coli* W3110 strain 9E4, which has the complete genotype *tonA ptr3*; *E. coli* W3110 strain 27C7 (ATCC 55,244), which has the complete genotype *tonA ptr3 phoA E15 (argF-lac)169 degP ompT kan^r*; *E. coli* W3110 strain 37D6, which has the complete genotype *tonA ptr3 phoA E15 (argF-lac)169 degP ompT rbs7 ilvG kan^r*; *E. coli* W3110 strain 40B4, which is strain 37D6 with a non-kanamycin resistant *degP* deletion mutation; and an *E. coli* strain having mutant periplasmic protease disclosed in U.S. Patent No. 4,946,783 issued 7 August 1990. Alternatively, *in vitro* methods of cloning, e.g., PCR or other nucleic acid polymerase reactions, are suitable.

In addition to prokaryotes, eukaryotic microbes such as filamentous fungi or yeast are suitable cloning or expression hosts for WISP-1 polypeptide-encoding vectors. *Saccharomyces cerevisiae* is a commonly used lower eukaryotic host microorganism. Others include *Schizosaccharomyces pombe* (Beach and Nurse, Nature, 290: 140 [1981]; EP 139,383 published 2 May 1985); *Kluyveromyces* hosts (U.S. Patent No. 4,943,529; Fleer et al., Bio/Technology, 9:968-975 (1991)) such as, e.g., *K. lactis* (MW98-8C, CBS683, CBS4574; Louvencourt et al., J. Bacteriol., 154(2):737-742 [1983]), *K. fragilis* (ATCC 12,424), *K. bulgaricus* (ATCC 16,045), *K. wickerhamii* (ATCC 24,178), *K. waltii* (ATCC 56,500), *K. drosophilae* (ATCC 36,906; Van den Berg et al., Bio/Technology, 8:135 (1990)), *K. thermotolerans*, and *K. marxianus*; *Yarrowia* (EP 402,226); *Pichia pastoris* (EP 183,070; Sreekrishna et al., J. Basic Microbiol., 28:265-278 [1988]); *Candida*; *Trichoderma reesia* (EP 244,234); *Neurospora crassa* (Case et al., Proc. Natl. Acad. Sci. USA, 76:5259-5263 [1979]); *Schwanniomyces* such as *Schwanniomyces occidentalis* (EP 394,538 published 31 October 1990); and filamentous fungi

such as, e.g., *Neurospora*, *Penicillium*, *Tolypocladium* (WO 91/00357 published 10 January 1991), and *Aspergillus* hosts such as *A. nidulans* (Ballance et al., Biochem. Biophys. Res. Commun., 112:284-289 [1983]; Tilburn et al., Gene, 26:205-221 [1983]; Yelton et al., Proc. Natl. Acad. Sci. USA, 81: 1470-1474 [1984]) and *A. niger* (Kelly and Hynes, EMBO J., 4:475-479 [1985]). Methylo-tropic yeasts are suitable herein and include, but are not limited to, yeast capable of growth on methanol selected from the genera consisting of *Hansenula*, *Candida*, *Kloeckera*, *Pichia*, *Saccharomyces*, *Torulopsis*, and *Rhodotorula*. A list of specific species that are exemplary of this class of yeasts may be found in C. Anthony, The Biochemistry of Methylo-trophs, 269 (1982).

Suitable host cells for the expression of glycosylated WISP-1 polypeptide are derived from multicellular organisms. Examples of invertebrate cells include insect cells such as *Drosophila* S2 and *Spodoptera* Sf9, as well as plant cells, such as cell cultures of cotton, corn, potato, soybean, petunia, tomato, and tobacco. Numerous baculoviral strains and variants and corresponding permissive insect host cells from hosts such as *Spodoptera frugiperda* (caterpillar), *Aedes aegypti* (mosquito), *Aedes albopictus* (mosquito), *Drosophila melanogaster* (fruitfly), and *Bombyx mori* have been identified. A variety of viral strains for transfection are publicly available, e.g., the L-1 variant of *Autographa californica* NPV and the Bm-5 strain of *Bombyx mori* NPV, and such viruses may be used as the virus herein according to the present invention, particularly for transfection of *Spodoptera frugiperda* cells.

However, interest has been greatest in vertebrate cells, and propagation of vertebrate cells in culture (tissue culture) has become a routine procedure. Examples of useful mammalian host cell lines are monkey kidney CV1 line transformed by SV40 (COS-7, ATCC CRL 1651); human embryonic kidney line (293 or 293 cells subcloned for growth in suspension culture, Graham et al., J. Gen Virol. 36:59 (1977)); baby hamster kidney cells (BHK, ATCC CCL 10); Chinese hamster ovary cells/-DHFR (CHO, Urlaub et al., Proc. Natl. Acad. Sci. USA 77:4216 (1980)); mouse sertoli cells (TM4, Mather, Biol. Reprod. 23:243-251 (1980)); monkey kidney cells (CV1 ATCC CCL 70); African green monkey kidney cells (VERO-76, ATCC CRL-1587); human cervical carcinoma cells (HELA, ATCC CCL 2); canine kidney cells (MDCK, ATCC CCL 34); buffalo rat liver cells (BRL 3A, ATCC CRL 1442); human lung cells (W138, ATCC CCL 75); human liver cells (Hep G2, HB 8065); mouse mammary tumor (MMT 060562, ATCC CCL51); TRI cells (Mather et al., Annals N.Y. Acad. Sci. 383:44-68 (1982)); MRC 5 cells; FS4 cells; and a human hepatoma line (Hep G2).

Host cells are transformed with the above-described expression or cloning vectors for WISP-1 polypeptide production and cultured in

conventional nutrient media modified as appropriate for inducing promoters, selecting transformants, or amplifying the genes encoding the desired sequences.

3. Selection and Use of a Replicable Vector

5 The nucleic acid (e.g., cDNA or genomic DNA) encoding WISP-1 polypeptide may be inserted into a replicable vector for cloning (amplification of the DNA) or for expression. Various vectors are publicly available. The vector may, for example, be in the form of a plasmid, cosmid, viral particle, or phage. The appropriate nucleic acid sequence
10 may be inserted into the vector by a variety of procedures. In general, DNA is inserted into an appropriate restriction endonuclease site(s) using techniques known in the art. Vector components generally include, but are not limited to, one or more of a signal sequence, an origin of replication, one or more marker genes, an enhancer element, a promoter, and a
15 transcription termination sequence. Construction of suitable vectors containing one or more of these components employs standard ligation techniques which are known to the skilled artisan.

The WISP-1 may be produced recombinantly not only directly, but also as a fusion polypeptide with a heterologous polypeptide, which may be a
20 signal sequence or other polypeptide having a specific cleavage site at the N-terminus of the mature protein or polypeptide. In general, the signal sequence may be a component of the vector, or it may be a part of the WISP-1 polypeptide-encoding DNA that is inserted into the vector. The signal sequence may be a prokaryotic signal sequence selected, for example, from
25 the group of the alkaline phosphatase, penicillinase, lpp, or heat-stable enterotoxin II leaders. For yeast secretion the signal sequence may be, e.g., the yeast invertase leader, alpha factor leader (including *Saccharomyces* and *Kluyveromyces* α -factor leaders, the latter described in U.S. Patent No. 5,010,182), or acid phosphatase leader, the *C. albicans*
30 glucoamylase leader (EP 362,179 published 4 April 1990), or the signal described in WO 90/13646 published 15 November 1990. In mammalian cell expression, mammalian signal sequences may be used to direct secretion of the protein, such as signal sequences from secreted polypeptides of the same or related species, as well as viral secretory leaders.

35 Both expression and cloning vectors contain a nucleic acid sequence that enables the vector to replicate in one or more selected host cells. Such sequences are well known for a variety of bacteria, yeast, and viruses. The origin of replication from the plasmid pBR322 is suitable for most Gram-negative bacteria, the 2 μ plasmid origin is suitable for yeast,
40 and various viral origins (SV40, polyoma, adenovirus, VSV or BPV) are useful for cloning vectors in mammalian cells.

Expression and cloning vectors will typically contain a selection gene, also termed a selectable marker. Typical selection genes encode proteins that (a) confer resistance to antibiotics or other toxins, e.g., ampicillin, neomycin, methotrexate, or tetracycline, (b) complement auxotrophic deficiencies, or (c) supply critical nutrients not available from complex media, e.g., the gene encoding D-alanine racemase for *Bacilli*.

An example of suitable selectable markers for mammalian cells are those that enable the identification of cells competent to take up the WISP-1 polypeptide-encoding nucleic acid, such as DHFR or thymidine kinase.

An appropriate host cell when wild-type DHFR is employed is the CHO cell line deficient in DHFR activity, prepared and propagated as described by Urlaub et al., Proc. Natl. Acad. Sci. USA, 77:4216 (1980). A suitable selection gene for use in yeast is the *trp1* gene present in the yeast plasmid YRp7 [Stinchcomb et al., Nature, 282:39 (1979); Kingsman et al., Gene, 7:141 (1979); Tschemper et al., Gene, 10:157 (1980)]. The *trp1* gene provides a selection marker for a mutant strain of yeast lacking the ability to grow in tryptophan, for example, ATCC No. 44076 or PEP4-1 [Jones, Genetics, 85:12 (1977)].

Expression and cloning vectors usually contain a promoter operably linked to the WISP-1 polypeptide-encoding nucleic acid sequence to direct mRNA synthesis. Promoters recognized by a variety of potential host cells are well known. Promoters suitable for use with prokaryotic hosts include the β -lactamase and lactose promoter systems [Chang et al., Nature, 275:615 (1978); Goeddel et al., Nature, 281:544 (1979)], alkaline phosphatase, a tryptophan (*trp*) promoter system [Goeddel, Nucleic Acids Res., 8:4057 (1980); EP 36,776], and hybrid promoters such as the *tac* promoter [deBoer et al., Proc. Natl. Acad. Sci. USA, 80:21-25 (1983)]. Promoters for use in bacterial systems also will contain a Shine-Dalgarno (S.D.) sequence operably linked to the DNA encoding WISP polypeptide.

Examples of suitable promoting sequences for use with yeast hosts include the promoters for 3-phosphoglycerate kinase [Hitzeman et al., J. Biol. Chem., 255:2073 (1980)] or other glycolytic enzymes [Hess et al., J. Adv. Enzyme Reg., 7:149 (1968); Holland, Biochemistry, 17:4900 (1978)], such as enolase, glyceraldehyde-3-phosphate dehydrogenase, hexokinase, pyruvate decarboxylase, phosphofructokinase, glucose-6-phosphate isomerase, 3-phosphoglycerate mutase, pyruvate kinase, triosephosphate isomerase, phosphoglucose isomerase, and glucokinase.

Other yeast promoters, which are inducible promoters having the additional advantage of transcription controlled by growth conditions, are the promoter regions for alcohol dehydrogenase 2, isocytochrome C, acid phosphatase, degradative enzymes associated with nitrogen metabolism, metallothionein, glyceraldehyde-3-phosphate dehydrogenase, and enzymes

responsible for maltose and galactose utilization. Suitable vectors and promoters for use in yeast expression are further described in EP 73,657.

WISP polypeptide transcription from vectors in mammalian host cells is controlled, for example, by promoters obtained from the genomes of viruses such as polyoma virus, fowlpox virus (UK 2,211,504 published 5 July 1989), adenovirus (such as Adenovirus 2), bovine papilloma virus, avian sarcoma virus, cytomegalovirus, a retrovirus, hepatitis-B virus and Simian Virus 40 (SV40), from heterologous mammalian promoters, e.g., the actin promoter or an immunoglobulin promoter, and from heat-shock promoters, provided such promoters are compatible with the host cell systems.

Transcription of a DNA encoding the WISP-1 polypeptide by higher eukaryotes may be increased by inserting an enhancer sequence into the vector. Enhancers are cis-acting elements of DNA, usually about from 10 to 300 bp, that act on a promoter to increase its transcription. Many enhancer sequences are now known from mammalian genes (globin, elastase, albumin, α -fetoprotein, and insulin). Typically, however, one will use an enhancer from a eukaryotic cell virus. Examples include the SV40 enhancer on the late side of the replication origin (bp 100-270), the cytomegalovirus early promoter enhancer, the polyoma enhancer on the late side of the replication origin, and adenovirus enhancers. The enhancer may be spliced into the vector at a position 5' or 3' to the WISP-1 polypeptide coding sequence, but is preferably located at a site 5' from the promoter.

Expression vectors used in eukaryotic host cells (yeast, fungi, insect, plant, animal, human, or nucleated cells from other multicellular organisms) will also contain sequences necessary for the termination of transcription and for stabilizing the mRNA. Such sequences are commonly available from the 5' and, occasionally 3', untranslated regions of eukaryotic or viral DNAs or cDNAs. These regions contain nucleotide segments transcribed as polyadenylated fragments in the untranslated portion of the mRNA encoding WISP-1 polypeptide.

Still other methods, vectors, and host cells suitable for adaptation to the synthesis of WISP polypeptide in recombinant vertebrate cell culture are described in Gething et al., Nature, 293:620-625 (1981); Mantei et al., Nature, 281:40-46 (1979); EP 117,060; and EP 117,058.

35 4. Culturing the Host Cells

The host cells used to produce the WISP polypeptide of this invention may be cultured in a variety of media. Commercially available media such as Ham's F10 (Sigma), Minimal Essential Medium (MEM); (Sigma), RPMI-1640 (Sigma), and Dulbecco's Modified Eagle's Medium ((DMEM), Sigma) are suitable for culturing the host cells. In addition, any of the media described in Ham et al., Meth. Enz. 58:44 (1979), Barnes et al., Anal. Biochem. 102:255 (1980), U.S. Pat. Nos. 4,767,704; 4,657,866; 4,927,762;

5,660,655; or 5,122,469; WO 90/03430; WO 87/00195; or U.S. Patent Re. 30,985 may be used as culture media for the host cells. Any of these media may be supplemented as necessary with hormones and/or other growth factors (such as insulin, transferrin, or epidermal growth factor), salts (such as sodium chloride, calcium, magnesium, and phosphate), buffers (such as HEPES), nucleotides (such as adenosine and thymidine), antibiotics (such as GENTAMYCIN™ drug), trace elements (defined as inorganic compounds usually present at final concentrations in the micromolar range), and glucose or an equivalent energy source. Any other necessary supplements may also be included at appropriate concentrations that would be known to those skilled in the art. The culture conditions, such as temperature, pH, and the like, are those previously used with the host cell selected for expression, and will be apparent to the ordinarily skilled artisan.

5. Detecting Gene Amplification/Expression

Gene amplification and/or expression may be measured in a sample directly, for example, by conventional Southern blotting, Northern blotting to quantitate the transcription of mRNA [Thomas, Proc. Natl. Acad. Sci. USA, 77:5201-5205 (1980)], dot blotting (DNA analysis), or *in situ* hybridization, using an appropriately labeled probe, based on the sequences provided herein. Alternatively, antibodies may be employed that can recognize specific duplexes, including DNA duplexes, RNA duplexes, and DNA-RNA hybrid duplexes or DNA-protein duplexes. The antibodies in turn may be labeled and the assay may be carried out where the duplex is bound to a surface, so that upon the formation of duplex on the surface, the presence of antibody bound to the duplex can be detected.

Gene expression, alternatively, may be measured by immunological methods, such as immunohistochemical staining of cells or tissue sections and assay of cell culture or body fluids, to quantitate directly the expression of gene product. Antibodies useful for immunohistochemical staining and/or assay of sample fluids may be either monoclonal or polyclonal, and may be prepared in any mammal. Conveniently, the antibodies may be prepared against a native sequence WISP polypeptide or against a synthetic peptide based on the DNA sequences provided herein or against exogenous sequence fused to WISP DNA and encoding a specific antibody epitope.

6. Purification of WISP Polypeptide

Forms of WISP polypeptide may be recovered from culture medium or from host cell lysates. If membrane-bound, it can be released from the membrane using a suitable detergent solution (e.g. Triton-X 100) or by enzymatic cleavage. Cells employed in expression of WISP-1 polypeptide can be disrupted by various physical or chemical means, such as freeze-thaw cycling, sonication, mechanical disruption, or cell lysing agents.

It may be desired to purify WISP-1 polypeptide from recombinant cell proteins or polypeptides. The following procedures are exemplary of suitable purification procedures: by fractionation on an ion-exchange column; ethanol precipitation; reverse phase HPLC; chromatography on silica or on a cation-exchange resin such as DEAE; chromatofocusing; SDS-PAGE; ammonium sulfate precipitation; gel filtration using, for example, Sephadex G-75; protein A Sepharose columns to remove contaminants such as IgG; and metal chelating columns to bind epitope-tagged forms of the WISP-1 polypeptide. Various methods of protein purification may be employed and such methods are known in the art and described for example in Deutscher, Methods in Enzymology, 182 (1990); Scopes, Protein Purification: Principles and Practice, Springer-Verlag, New York (1982). The purification step(s) selected will depend, for example, on the nature of the production process used and the particular WISP-1 polypeptide produced.

Soluble forms of WISP-1 may be employed as antagonists in the methods of the invention. Such soluble forms of WISP-1 may comprise modifications, as described below (such as by fusing to an immunoglobulin, epitope tag or leucine zipper). Immunoadhesin molecules are further contemplated for use in the methods herein. WISP-1 immunoadhesins may comprise various forms of WISP-1, such as the full length polypeptide as well as soluble forms of the WISP-1 or a fragment thereof. In particular embodiments, the molecule may comprise a fusion of the WISP-1 polypeptide with an immunoglobulin or a particular region of an immunoglobulin. For a bivalent form of the immunoadhesin, such a fusion could be to the Fc region of an IgG molecule. The Ig fusions preferably include the substitution of a soluble (transmembrane domain deleted or inactivated) form of the polypeptide in place of at least one variable region within an Ig molecule. In a particularly preferred embodiment, the immunoglobulin fusion includes the hinge, CH2 and CH3, or the hinge, CH1, CH2 and CH3 regions of an IgG1 molecule. For the production of immunoglobulin fusions, see also US Patent No. 5,428,130 issued June 27, 1995 and Chamow et al., TIBTECH, 14:52-60 (1996).

The simplest and most straightforward immunoadhesin design combines the binding domain(s) of the adhesin (e.g. the WISP-1) with the Fc region of an immunoglobulin heavy chain. Ordinarily, when preparing the immunoadhesins of the present invention, nucleic acid encoding the binding domain of the adhesin will be fused C-terminally to nucleic acid encoding the N-terminus of an immunoglobulin constant domain sequence, however N-terminal fusions are also possible.

Typically, in such fusions the encoded chimeric polypeptide will retain at least functionally active hinge, CH2 and CH3 domains of the constant region of an immunoglobulin heavy chain. Fusions are also made to

the C-terminus of the Fc portion of a constant domain, or immediately N-terminal to the C_H1 of the heavy chain or the corresponding region of the light chain. The precise site at which the fusion is made is not critical; particular sites are well known and may be selected in order to optimize the biological activity, secretion, or binding characteristics of the immunoadhesin.

In a preferred embodiment, the adhesin sequence is fused to the N-terminus of the Fc region of immunoglobulin G₁ (IgG₁). It is possible to fuse the entire heavy chain constant region to the adhesin sequence. However, more preferably, a sequence beginning in the hinge region just upstream of the papain cleavage site which defines IgG Fc chemically (i.e. residue 216, taking the first residue of heavy chain constant region to be 114), or analogous sites of other immunoglobulins is used in the fusion. In a particularly preferred embodiment, the adhesin amino acid sequence is fused to (a) the hinge region and C_H2 and C_H3 or (b) the C_H1, hinge, C_H2 and C_H3 domains, of an IgG heavy chain.

For bispecific immunoadhesins, the immunoadhesins are assembled as multimers, and particularly as heterodimers or heterotetramers. Generally, these assembled immunoglobulins will have known unit structures. A basic four chain structural unit is the form in which IgG, IgD, and IgE exist. A four chain unit is repeated in the higher molecular weight immunoglobulins; IgM generally exists as a pentamer of four basic units held together by disulfide bonds. IgA globulin, and occasionally IgG globulin, may also exist in multimeric form in serum. In the case of multimer, each of the four units may be the same or different.

Various exemplary assembled immunoadhesins within the scope herein are schematically diagrammed below:

- (a) AC_L-AC_L;
- (b) AC_H-(AC_H, AC_L-AC_H, AC_L-V_HC_H, or V_LC_L-AC_H);
- (c) AC_L-AC_H-(AC_L-AC_H, AC_L-V_HC_H, V_LC_L-AC_H, or V_LC_L-V_HC_H);
- (d) AC_L-V_HC_H-(AC_H, or AC_L-V_HC_H, or V_LC_L-AC_H);
- (e) V_LC_L-AC_H-(AC_L-V_HC_H, or V_LC_L-AC_H); and
- (f) (A-Y)_n-(V_LC_L-V_HC_H)₂,

wherein each A represents identical or different adhesin amino acid sequences;

V_L is an immunoglobulin light chain variable domain;
V_H is an immunoglobulin heavy chain variable domain;
C_L is an immunoglobulin light chain constant domain;
C_H is an immunoglobulin heavy chain constant domain;
n is an integer greater than 1;
Y designates the residue of a covalent cross-linking agent.

In the interests of brevity, the foregoing structures only show key features; they do not indicate joining (J) or other domains of the immunoglobulins, nor are disulfide bonds shown. However, where such domains are required for binding activity, they shall be constructed to be present in the ordinary locations which they occupy in the immunoglobulin molecules.

Alternatively, the adhesin sequences can be inserted between immunoglobulin heavy chain and light chain sequences, such that an immunoglobulin comprising a chimeric heavy chain is obtained. In this embodiment, the adhesin sequences are fused to the 3' end of an immunoglobulin heavy chain in each arm of an immunoglobulin, either between the hinge and the C_H2 domain, or between the C_H2 and C_H3 domains. Similar constructs have been reported by Hoogenboom et al., Mol. Immunol., 28:1027-1037 (1991).

Although the presence of an immunoglobulin light chain is not required in the immunoadhesins of the present invention, an immunoglobulin light chain might be present either covalently associated to an adhesin-immunoglobulin heavy chain fusion polypeptide, or directly fused to the adhesin. In the former case, DNA encoding an immunoglobulin light chain is typically coexpressed with the DNA encoding the adhesin-immunoglobulin heavy chain fusion protein. Upon secretion, the hybrid heavy chain and the light chain will be covalently associated to provide an immunoglobulin-like structure comprising two disulfide-linked immunoglobulin heavy chain-light chain pairs. Methods suitable for the preparation of such structures are, for example, disclosed in U.S. Patent No. 4,816,567, issued 28 March 1989.

Immunoadhesins are most conveniently constructed by fusing the cDNA sequence encoding the adhesin portion in-frame to an immunoglobulin cDNA sequence. However, fusion to genomic immunoglobulin fragments can also be used (see, e.g. Aruffo et al., Cell, 61:1303-1313 (1990); and Stamenkovic et al., Cell, 66:1133-1144 (1991)). The latter type of fusion requires the presence of Ig regulatory sequences for expression. cDNAs encoding IgG heavy-chain constant regions can be isolated based on published sequences from cDNA libraries derived from spleen or peripheral blood lymphocytes, by hybridization or by polymerase chain reaction (PCR) techniques. The cDNAs encoding the "adhesin" and the immunoglobulin parts of the immunoadhesin are inserted in tandem into a plasmid vector that directs efficient expression in the chosen host cells.

In another embodiment, the WISP-1 or WISP-1 antagonist may be covalently modified by linking the receptor polypeptide to one of a variety of nonproteinaceous polymers, e.g., polyethylene glycol (PEG), polypropylene glycol, or polyoxyalkylenes, in the manner set forth in U.S. Patent Nos. 4,640,835; 4,496,689; 4,301,144; 4,670,417; 4,791,192 or

Leucine zipper forms of these molecules are also contemplated by the invention. "Leucine zipper" is a term in the art used to refer to a leucine rich sequence that enhances, promotes, or drives dimerization or trimerization of its fusion partner (e.g., the sequence or molecule to which the leucine zipper is fused or linked to). Various leucine zipper polypeptides have been described in the art. See, e.g., Landschulz et al., Science, 240:1759 (1988); US Patent 5,716,805; WO 94/10308; Hoppe et al., FEBS Letters, 344:1991 (1994); Maniatis et al., Nature, 341:24 (1989). Those skilled in the art will appreciate that a leucine zipper sequence may be fused at either the 5' or 3' end of the WISP-1 or WISP-1 antagonist molecule.

The WISP-1 polypeptides of the present invention may also be modified in a way to form chimeric molecules by fusing the polypeptide to another, heterologous polypeptide or amino acid sequence. Preferably, such heterologous polypeptide or amino acid sequence is one which acts to oligomerize the chimeric molecule. In one embodiment, such a chimeric molecule comprises a fusion of the WISP-1 polypeptide with a tag polypeptide which provides an epitope to which an anti-tag antibody can selectively bind. The epitope tag is generally placed at the amino- or carboxyl- terminus of the polypeptide. The presence of such epitope-tagged forms of the polypeptide can be detected using an antibody against the tag polypeptide. Also, provision of the epitope tag enables the polypeptide to be readily purified by affinity purification using an anti-tag antibody or another type of affinity matrix that binds to the epitope tag. Various tag polypeptides and their respective antibodies are well known in the art. Examples include poly-histidine (poly-his) or poly-histidine-glycine (poly-his-gly) tags; the flu HA tag polypeptide and its antibody 12CA5 [Field et al., Mol. Cell. Biol., 8:2159-2165 (1988)]; the c-myc tag and the 8F9, 3C7, 6E10, G4, B7 and 9E10 antibodies thereto [Evan et al., Molecular and Cellular Biology, 5:3610-3616 (1985)]; and the Herpes Simplex virus glycoprotein D (gD) tag and its antibody [Paborsky et al., Protein Engineering, 3(6):547-553 (1990)]. Other tag polypeptides include the Flag-peptide [Hopp et al., BioTechnology, 6:1204-1210 (1988)]; the KT3 epitope peptide [Martin et al., Science, 255:192-194 (1992)]; an α -tubulin epitope peptide [Skinner et al., J. Biol. Chem., 266:15163-15166 (1991)]; and the T7 gene 10 protein peptide tag [Lutz-Freyermuth et al., Proc. Natl. Acad. Sci. USA, 87:6393-6397 (1990)].

It is contemplated that anti-WISP-1 antibodies may also be employed in the presently disclosed methods. Examples of such molecules include

Neutralizing or blocking antibodies which can reduce cancer cell growth, metastasis or motility. The anti-WISP-1 may be monoclonal antibodies.

Monoclonal antibodies may be prepared using hybridoma methods, such as those described by Kohler and Milstein, Nature, 256:495 (1975). In a hybridoma method, a mouse, hamster, or other appropriate host animal, is typically immunized with an immunizing agent to elicit lymphocytes that produce or are capable of producing antibodies that will specifically bind to the immunizing agent. Alternatively, the lymphocytes may be immunized in vitro.

The immunizing agent will typically include a WISP-1 polypeptide or a fusion protein thereof, such as a WISP-1-IgG fusion protein.

Generally, either peripheral blood lymphocytes ("PBLs") are used if cells of human origin are desired, or spleen cells or lymph node cells are used if non-human mammalian sources are desired. The lymphocytes are then fused with an immortalized cell line using a suitable fusing agent, such as polyethylene glycol, to form a hybridoma cell [Goding, Monoclonal Antibodies: Principles and Practice, Academic Press, (1986) pp. 59-103]. Immortalized cell lines are usually transformed mammalian cells, particularly myeloma cells of rodent, bovine and human origin. Usually, rat or mouse myeloma cell lines are employed. The hybridoma cells may be cultured in a suitable culture medium that preferably contains one or more substances that inhibit the growth or survival of the unfused, immortalized cells. For example, if the parental cells lack the enzyme hypoxanthine guanine phosphoribosyl transferase (HGPRT or HPRT), the culture medium for the hybridomas typically will include hypoxanthine, aminopterin, and thymidine ("HAT medium"), which substances prevent the growth of HGPRT-deficient cells.

Preferred immortalized cell lines are those that fuse efficiently, support stable high level expression of antibody by the selected antibody-producing cells, and are sensitive to a medium such as HAT medium. More preferred immortalized cell lines are murine myeloma lines, which can be obtained, for instance, from the Salk Institute Cell Distribution Center, San Diego, California and the American Type Culture Collection, Manassas, Virginia. Human myeloma and mouse-human heteromyeloma cell lines also have been described for the production of human monoclonal antibodies [Kozbor, J. Immunol., 133:3001 (1984); Brodeur et al., Monoclonal Antibody Production Techniques and Applications, Marcel Dekker, Inc., New York, (1987) pp. 51-63].

The culture medium in which the hybridoma cells are cultured can then be assayed for the presence of monoclonal antibodies directed against WISP-1. Preferably, the binding specificity of monoclonal antibodies produced by the hybridoma cells is determined by immunoprecipitation or by an in vitro binding assay, such as radioimmunoassay (RIA) or enzyme-linked

immunoabsorbent assay (ELISA). Such techniques and assays are known in the art. The binding affinity of the monoclonal antibody can, for example, be determined by the Scatchard analysis of Munson and Pollard, Anal. Biochem., 107:220 (1980).

5 After the desired hybridoma cells are identified, the clones may be subcloned by limiting dilution procedures and grown by standard methods [Goding, supra]. Suitable culture media for this purpose include, for example, Dulbecco's Modified Eagle's Medium or RPMI-1640 medium. Alternatively, the hybridoma cells may be grown *in vivo* as ascites in a
10 mammal.

The monoclonal antibodies secreted by the subclones may be isolated or purified from the culture medium or ascites fluid by conventional immunoglobulin purification procedures such as, for example, protein A-Sepharose, hydroxylapatite chromatography, gel electrophoresis, dialysis, or
15 affinity chromatography.

The monoclonal antibodies may also be made by recombinant DNA methods, such as those described in U.S. Patent No. 4,816,567. DNA encoding the monoclonal antibodies is readily isolated and sequenced using conventional procedures (e.g., by using oligonucleotide probes that are capable of
20 binding specifically to genes encoding the heavy and light chains of the monoclonal antibodies). The hybridoma cells serve as a preferred source of such DNA. Once isolated, the DNA may be placed into expression vectors, which are then transfected into host cells such as *E. coli* cells, simian COS cells, Chinese hamster ovary (CHO) cells, or myeloma cells that do not
25 otherwise produce immunoglobulin protein, to obtain the synthesis of monoclonal antibodies in the recombinant host cells. The DNA also may be modified, for example, by substituting the coding sequence for human heavy and light chain constant domains in place of the homologous murine sequences, Morrison, et al., Proc. Nat. Acad. Sci. 81, 6851 (1984), or by
30 covalently joining to the immunoglobulin coding sequence all or part of the coding sequence for a non-immunoglobulin polypeptide.

Typically such non-immunoglobulin polypeptides are substituted for the constant domains of an antibody of the invention, or they are substituted for the variable domains of one antigen-combining site of an
35 antibody of the invention to create a chimeric bivalent antibody comprising one antigen-combining site having specificity for WISP-1 and another antigen-combining site having specificity for a different antigen.

Chimeric or hybrid antibodies also may be prepared in vitro using known methods in synthetic protein chemistry, including those involving
40 crosslinking agents. For example, immunotoxins may be constructed using a disulfide exchange reaction or by forming a thioether bond. Examples of

Suitable reagents for this purpose include iminothiolate and methyl-4-mercaptobutyrimidate.

Single chain Fv fragments may also be produced, such as described in Iliades et al., FEBS Letters, 409:437-441 (1997). Coupling of such single chain fragments using various linkers is described in Kortt et al., Protein Engineering, 10:423-433 (1997). A variety of techniques for the recombinant production and manipulation of antibodies are well known in the art. Illustrative examples of such techniques that are typically utilized by skilled artisans are described in greater detail below.

10 (i) *Humanized antibodies*

Generally, a humanized antibody has one or more amino acid residues introduced into it from a non-human source. These non-human amino acid residues are often referred to as "import" residues, which are typically taken from an "import" variable domain. Humanization can be essentially performed following the method of Winter and co-workers [Jones et al., Nature, 321:522-525 (1986); Riechmann et al., Nature, 332:323-327 (1988); Verhoeyen et al., Science, 239:1534-1536 (1988)], by substituting rodent CDRs or CDR sequences for the corresponding sequences of a human antibody.

Accordingly, such "humanized" antibodies are chimeric antibodies wherein substantially less than an intact human variable domain has been substituted by the corresponding sequence from a non-human species. In practice, humanized antibodies are typically human antibodies in which some CDR residues and possibly some FR residues are substituted by residues from analogous sites in rodent antibodies.

25 It is important that antibodies be humanized with retention of high affinity for the antigen and other favorable biological properties. To achieve this goal, according to a preferred method, humanized antibodies are prepared by a process of analysis of the parental sequences and various conceptual humanized products using three dimensional models of the parental and humanized sequences. Three dimensional immunoglobulin models are commonly available and are familiar to those skilled in the art. Computer programs are available which illustrate and display probable three-dimensional conformational structures of selected candidate immunoglobulin sequences. Inspection of these displays permits analysis of the likely role of the residues in the functioning of the candidate immunoglobulin sequence, i.e. the analysis of residues that influence the ability of the candidate immunoglobulin to bind its antigen. In this way, FR residues can be selected and combined from the consensus and import sequence so that the desired antibody characteristic, such as increased affinity for the target antigen(s), is achieved. In general, the CDR residues are directly and most substantially involved in influencing antigen binding.

(ii) Human antibodies

Human monoclonal antibodies can be made by the hybridoma method.

Human myeloma and mouse-human heteromyeloma cell lines for the production of human monoclonal antibodies have been described, for example, by Kozbor, J. Immunol. 133, 3001 (1984), and Brodeur, et al., Monoclonal Antibody Production Techniques and Applications, pp.51-63 (Marcel Dekker, Inc., New York, 1987).

It is now possible to produce transgenic animals (e.g. mice) that are capable, upon immunization, of producing a repertoire of human antibodies in the absence of endogenous immunoglobulin production. For example, it has been described that the homozygous deletion of the antibody heavy chain joining region (J_H) gene in chimeric and germ-line mutant mice results in complete inhibition of endogenous antibody production. Transfer of the human germ-line immunoglobulin gene array in such germ-line mutant mice will result in the production of human antibodies upon antigen challenge.

See, e.g. Jakobovits et al., Proc. Natl. Acad. Sci. USA 90, 2551-255 (1993); Jakobovits et al., Nature 362, 255-258 (1993).

Mendez et al. (Nature Genetics 15: 146-156 [1997]) have further improved the technology and have generated a line of transgenic mice designated as "Xenomouse II" that, when challenged with an antigen, generates high affinity fully human antibodies. This was achieved by germ-line integration of megabase human heavy chain and light chain loci into mice with deletion into endogenous J_H segment as described above. The Xenomouse II harbors 1,020 kb of human heavy chain locus containing approximately 66 V_H genes, complete D_H and J_H regions and three different constant regions (μ , δ and χ), and also harbors 800 kb of human κ locus containing 32 V_k genes, J_k segments and C_k genes. The antibodies produced in these mice closely resemble that seen in humans in all respects, including gene rearrangement, assembly, and repertoire. The human antibodies are preferentially expressed over endogenous antibodies due to deletion in endogenous J_H segment that prevents gene rearrangement in the murine locus.

Alternatively, the phage display technology (McCafferty et al., Nature 348, 552-553 [1990]) can be used to produce human antibodies and antibody fragments *in vitro*, from immunoglobulin variable (V) domain gene repertoires from unimmunized donors. According to this technique, antibody V domain genes are cloned in-frame into either a major or minor coat protein gene of a filamentous bacteriophage, such as M13 or fd, and displayed as functional antibody fragments on the surface of the phage particle. Because the filamentous particle contains a single-stranded DNA copy of the phage genome, selections based on the functional properties of the antibody also result in selection of the gene encoding the antibody

forms of antibodies. Those skilled in the art may construct such dimers or multivalent forms by techniques known in the art. Methods for preparing monovalent antibodies are also well known in the art. For example, one method involves recombinant expression of immunoglobulin light chain and modified heavy chain. The heavy chain is truncated generally at any point in the Fc region so as to prevent heavy chain crosslinking. Alternatively, the relevant cysteine residues are substituted with another amino acid residue or are deleted so as to prevent crosslinking.

(iii) *Bispecific antibodies*

Bispecific antibodies are monoclonal, preferably human or humanized, antibodies that have binding specificities for at least two different antigens. In the present case, one of the binding specificities is for WISP-1. For example, bispecific antibodies specifically binding WISP-1 or WISP-1 variants and another CNN family member (e.g., WISP-2, WISP-3, CTGF, Cyr61, or Nov) or other molecules such as CD44 are within the scope of the present invention.

Methods for making bispecific antibodies are known in the art. Traditionally, the recombinant production of bispecific antibodies is based on the coexpression of two immunoglobulin heavy chain-light chain pairs, where the two heavy chains have different specificities (Millstein and Cuello, *Nature* 305, 537-539 (1983)). Because of the random assortment of immunoglobulin heavy and light chains, these hybridomas (quadromas) produce a potential mixture of 10 different antibody molecules, of which only one has the correct bispecific structure. The purification of the correct molecule, which is usually done by affinity chromatography steps, is rather cumbersome, and the product yields are low. Similar procedures are disclosed in PCT application publication No. WO 93/08829 (published 13 May 1993), and in Traunecker et al., *EMBO* 10, 3655-3659 (1991).

According to a different and more preferred approach, antibody variable domains with the desired binding specificities (antibody-antigen combining sites) are fused to immunoglobulin constant domain sequences. The fusion preferably is with an immunoglobulin heavy chain constant domain, comprising at least part of the hinge, CH2 and CH3 regions. It is preferred to have the first heavy chain constant region (CH1) containing the site necessary for light chain binding, present in at least one of the fusions. DNAs encoding the immunoglobulin heavy chain fusions and, if desired, the immunoglobulin light chain, are inserted into separate expression vectors, and are cotransfected into a suitable host organism. This provides for great flexibility in adjusting the mutual proportions of the three polypeptide fragments in embodiments when unequal ratios of the three polypeptide chains used in the construction provide the optimum yields. It is, however, possible to insert the coding sequences for two or

all three polypeptide chains in one expression vector when the expression of at least two polypeptide chains in equal ratios results in high yields or when the ratios are of no particular significance. In a preferred embodiment of this approach, the bispecific antibodies are composed of a hybrid immunoglobulin heavy chain with a first binding specificity in one arm, and a hybrid immunoglobulin heavy chain-light chain pair (providing a second binding specificity) in the other arm. It was found that this asymmetric structure facilitates the separation of the desired bispecific compound from unwanted immunoglobulin chain combinations, as the presence of an immunoglobulin light chain in only one half of the bispecific molecule provides for a facile way of separation. This approach is disclosed in PCT Publication No. WO 94/04690, published on March 3, 1994.

For further details of generating bispecific antibodies see, for example, Suresh et al., Methods in Enzymology 121, 210 (1986).

(iv) *Heteroconjugate antibodies*

Heteroconjugate antibodies are also within the scope of the present invention. Heteroconjugate antibodies are composed of two covalently joined antibodies. Such antibodies have, for example, been proposed to target immune system cells to unwanted cells (U.S. Patent No. 4,676,980), and for treatment of HIV infection (PCT application publication Nos. WO 91/00360 and WO 92/200373; EP 03089). Heteroconjugate antibodies may be made using any convenient cross-linking methods. Suitable cross-linking agents are well known in the art, and are disclosed in U.S. Patent No. 4,676,980, along with a number of cross-linking techniques.

(v) *Antibody fragments*

In certain embodiments, the anti-WISP-1 antibody (including murine, human and humanized antibodies, and antibody variants) is an antibody fragment. Various techniques have been developed for the production of antibody fragments. Traditionally, these fragments were derived via proteolytic digestion of intact antibodies (see, e.g., Morimoto et al., *J. Biochem. Biophys. Methods* 24:107-117 (1992) and Brennan et al., *Science* 229:81 (1985)). However, these fragments can now be produced directly by recombinant host cells. For example, Fab'-SH fragments can be directly recovered from *E. coli* and chemically coupled to form F(ab')₂ fragments (Carter et al., *Bio/Technology* 10:163-167 (1992)). In another embodiment, the F(ab')₂ is formed using the leucine zipper GCN4 to promote assembly of the F(ab')₂ molecule. According to another approach, Fv, Fab or F(ab')₂ fragments can be isolated directly from recombinant host cell culture. A variety of techniques for the production of antibody fragments will be apparent to the skilled practitioner. For instance, digestion can be performed using papain. Examples of papain digestion are described in WO 94/29348 published 12/22/94 and U.S. Patent No. 4,342,566. Papain digestion

of antibodies typically produces two identical antigen binding fragments, called Fab fragments, each with a single antigen binding site, and a residual Fc fragment. Pepsin treatment yields an $F(ab')_2$ fragment that has two antigen combining sites and is still capable of cross-linking antigen.

5 The Fab fragments produced in the antibody digestion also contain the constant domains of the light chain and the first constant domain (CH_1) of the heavy chain. Fab' fragments differ from Fab fragments by the addition of a few residues at the carboxy terminus of the heavy chain CH_1 domain including one or more cysteines from the antibody hinge region. Fab'-SH is
10 the designation herein for Fab' in which the cysteine residue(s) of the constant domains bear a free thiol group. $F(ab')_2$ antibody fragments originally were produced as pairs of Fab' fragments which have hinge cysteines between them. Other chemical couplings of antibody fragments are also known.

15 Antibodies are glycosylated at conserved positions in their constant regions (Jefferis and Lund, Chem. Immunol. 65:111-128 [1997]; Wright and Morrison, TibTECH 15:26-32 [1997]). The oligosaccharide side chains of the immunoglobulins affect the protein's function (Boyd et al., Mol. Immunol. 32:1311-1318 [1996]; Wittwe and Howard, Biochem. 29:4175-4180
20 [1990]), and the intramolecular interaction between portions of the glycoprotein which can affect the conformation and presented three-dimensional surface of the glycoprotein (Hefferis and Lund, supra; Wyss and Wagner, Current Opin. Biotech. 7:409-416 [1996]). Oligosaccharides may also serve to target a given glycoprotein to certain molecules based upon specific recognition structures. For example, it has been reported
25 that in agalactosylated IgG, the oligosaccharide moiety 'flips' out of the inter- CH_2 space and terminal N-acetylglucosamine residues become available to bind mannose binding protein (Malhotra et al., Nature Med. 1:237-243 [1995]). Removal by glycopeptidase of the oligosaccharides from CAMPATH-
30 1H (a recombinant humanized murine monoclonal IgG1 antibody which recognizes the CDw52 antigen of human lymphocytes) produced in Chinese Hamster Ovary (CHO) cells resulted in a complete reduction in complement mediated lysis (CMCL) (Boyd et al., Mol. Immunol. 32:1311-1318 [1996]), while selective removal of sialic acid residues using neuraminidase
35 resulted in no loss of DMCL. Glycosylation of antibodies has also been reported to affect antibody-dependent cellular cytotoxicity (ADCC). In particular, CHO cells with tetracycline-regulated expression of $\beta(1,4)$ -N-acetylglucosaminyltransferase III (GnTIII), a glycosyltransferase catalyzing formation of bisecting GlcNAc, was reported to have improved
40 ADCC activity (Umana et al., Mature Biotech. 17:176-180 [1999]).

Glycosylation variants of antibodies are variants in which the glycosylation pattern of an antibody is altered. By altering is meant

deleting one or more carbohydrate moieties found in the antibody, adding one or more carbohydrate moieties to the antibody, changing the composition of glycosylation (glycosylation pattern), the extent of glycosylation, etc. Glycosylation variants may, for example, be prepared by removing, changing and/or adding one or more glycosylation sites in the nucleic acid sequence encoding the antibody.

Glycosylation of antibodies is typically either N-linked or O-linked. N-linked refers to the attachment of the carbohydrate moiety to the side chain of an asparagine residue. The tripeptide sequences asparagine-X-serine and asparagine-X-threonine, where X is any amino acid except proline, are the recognition sequences for enzymatic attachment of the carbohydrate moiety to the asparagine side chain. Thus, the presence of either of these tripeptide sequences in a polypeptide creates a potential glycosylation site. O-linked glycosylation refers to the attachment of one of the sugars N-acetylgalactosamine, galactose, or xylose to a hydroxyamino acid, most commonly serine or threonine, although 5-hydroxyproline or 5-hydroxylysine may also be used.

Addition of glycosylation sites to the antibody is conveniently accomplished by altering the amino acid sequence such that it contains one or more of the above-described tripeptide sequences (for N-linked glycosylation sites). The alteration may also be made by the addition of, or substitution by, one or more serine or threonine residues to the sequence of the original antibody (for O-linked glycosylation sites).

The glycosylation (including glycosylation pattern) of antibodies may also be altered without altering the underlying nucleotide sequence. Glycosylation largely depends on the host cell used to express the antibody. Since the cell type used for expression of recombinant glycoproteins, e.g. antibodies, as potential therapeutics is rarely the native cell, significant variations in the glycosylation pattern of the antibodies can be expected (see, e.g. Hse et al., J. Biol. Chem. 272:9062-9070 [1997]). In addition to the choice of host cells, factors which affect glycosylation during recombinant production of antibodies include growth mode, media formulation, culture density, oxygenation, pH, purification schemes and the like. Various methods have been proposed to alter the glycosylation pattern achieved in a particular host organism including introducing or overexpressing certain enzymes involved in oligosaccharide production (U. S. Patent Nos. 5,047,335; 5,510,261 and 5,278,299). Glycosylation, or certain types of glycosylation, can be enzymatically removed from the glycoprotein, for example using endoglycosidase H (Endo H). In addition, the recombinant host cell can be genetically engineered, e.g. make defective in processing certain types of polysaccharides. These and similar techniques are well known in the art.

Massey, *Nature* 328: 457-458 (1987)). Antibody-abzyme conjugates can be prepared as described herein for delivery of the abzyme to a tumor cell population.

The enzymes can be covalently bound to the antibodies by techniques well known in the art such as the use of heterobifunctional crosslinking reagents. Alternatively, fusion proteins comprising at least the antigen binding region of an antibody of the invention linked to at least a functionally active portion of an enzyme of the invention can be constructed using recombinant DNA techniques well known in the art (see, e.g., Neuberger et al., *Nature*, 312: 604-608 (1984)).

Further antibody modifications are contemplated. For example, the antibody may be linked to one of a variety of nonproteinaceous polymers, e.g., polyethylene glycol, polypropylene glycol, polyoxyalkylenes, or copolymers of polyethylene glycol and polypropylene glycol, or other molecules such as polyglutamate. The antibody also may be entrapped in microcapsules prepared, for example, by coacervation techniques or by interfacial polymerization (for example, hydroxymethylcellulose or gelatin-microcapsules and poly-(methylmethacrylate) microcapsules, respectively), in colloidal drug delivery systems (for example, liposomes, albumin microspheres, microemulsions, nano-particles and nanocapsules), or in macroemulsions. Such techniques are disclosed in Remington's *Pharmaceutical Sciences*, 16th edition, Oslo, A., Ed., (1980). To increase the serum half life of the antibody, one may incorporate a salvage receptor binding epitope into the antibody (especially an antibody fragment) as described in U.S. Patent 5,739,277, for example. As used herein, the term "salvage receptor binding epitope" refers to an epitope of the Fc region of an IgG molecule (e.g., IgG₁, IgG₂, IgG₃, or IgG₄) that is responsible for increasing the *in vivo* serum half-life of the IgG molecule.

FORMULATIONS

The WISP-1 antagonists described herein, are optionally employed in a carrier. Suitable carriers and their formulations are described in Remington's Pharmaceutical Sciences, 16th ed., 1980, Mack Publishing Co., edited by Osol et al. Typically, an appropriate amount of a pharmaceutically-acceptable salt is used in the carrier to render the formulation isotonic. Examples of the carrier include saline, Ringer's solution and dextrose solution. The pH of the carrier is preferably from about 5 to about 8, and more preferably from about 7.4 to about 7.8. It will be apparent to those persons skilled in the art that certain carriers may be more preferable depending upon, for instance, the route of administration and concentration of active agent being administered. The carrier may be in the form of a lyophilized formulation or aqueous solution.

Acceptable carriers, excipients, or stabilizers are preferably nontoxic to cells and/or recipients at the dosages and concentrations employed, and include buffers such as phosphate, citrate, and other organic acids; antioxidants including ascorbic acid and methionine; preservatives (such as octadecyldimethylbenzyl ammonium chloride; hexamethonium chloride; benzalkonium chloride, benzethonium chloride; phenol, butyl or benzyl alcohol; alkyl parabens such as methyl or propyl paraben; catechol; resorcinol; cyclohexanol; 3-pentanol; and m-cresol); low molecular weight (less than about 10 residues) polypeptides; proteins, such as serum albumin, gelatin, or immunoglobulins; hydrophilic polymers such as polyvinylpyrrolidone; amino acids such as glycine, glutamine, asparagine, histidine, arginine, or lysine; monosaccharides, disaccharides, and other carbohydrates including glucose, mannose, or dextrans; chelating agents such as EDTA; sugars such as sucrose, mannitol, trehalose or sorbitol; salt-forming counter-ions such as sodium; and/or non-ionic surfactants such as TWEEN™, PLURONICS™ or polyethylene glycol (PEG).

The formulation may also contain more than one active compound as necessary for the particular indication being treated, preferably those with complementary activities that do not adversely affect each other.

The antagonists described herein may also be entrapped in microcapsules prepared, for example, by coacervation techniques or by interfacial polymerization, for example, hydroxymethylcellulose or gelatin-microcapsules and poly-(methylmethacrylate) microcapsules, respectively, in colloidal drug delivery systems (for example, liposomes, albumin microspheres, microemulsions, nano-particles and nanocapsules) or in macroemulsions. Such techniques are disclosed in Remington's Pharmaceutical Sciences 16th edition, Osol, A. Ed. (1980).

The formulations to be used for in vivo administration should be sterile. This is readily accomplished by filtration through sterile filtration membranes.

Sustained-release preparations may be prepared. Suitable examples of sustained-release preparations include semipermeable matrices of solid hydrophobic polymers containing the active agent, which matrices are in the form of shaped articles, e.g. films, or microcapsules. Examples of sustained-release matrices include polyesters, hydrogels (for example, poly(2-hydroxyethyl-methacrylate), or poly(vinylalcohol)), polylactides (U.S. Pat. No. 3,773,919), copolymers of L-glutamic acid and γ ethyl-L-glutamate, non-degradable ethylene-vinyl acetate, degradable lactic acid-glycolic acid copolymers such as the LUPRON DEPOT™ (injectable microspheres composed of lactic acid-glycolic acid copolymer and leuprolide acetate), and poly-D-(-)-3-hydroxybutyric acid. While polymers such as ethylene-vinyl acetate and lactic acid-glycolic acid enable release of

molecules for over 100 days, certain hydrogels release proteins for shorter time periods.

MODES OF THERAPY

MODES OF THERAPY

The molecules described herein are useful in treating various pathological conditions, such as cancer. These conditions can be treated by administration of one or more antagonists described herein. Diagnosis in mammals of the various pathological conditions described herein can be made by the skilled practitioner. Diagnostic techniques are available in the art which allow, e.g., for the diagnosis or detection of cancer or immune related disease in a mammal. For instance, cancers may be identified through techniques, including but not limited to, palpation, blood analysis, x-ray, NMR and the like.

The antagonist(s) can be administered in accord with known methods, such as intravenous administration as a bolus or by continuous infusion over a period of time, by intramuscular, intraperitoneal, intracerebrospinal, subcutaneous, intra-articular, intrasynovial, intrathecal, oral, topical, or inhalation routes. Optionally, administration may be performed through mini-pump infusion using various commercially available devices. The antagonists may also be employed using gene therapy techniques which have been described in the art.

Effective dosages and schedules for administering antagonists may be determined empirically, and making such determinations is within the skill in the art. Single or multiple dosages may be employed. It is presently believed that an effective dosage or amount of antagonist used alone may range from about 1 µg/kg to about 100 mg/kg of body weight or more per day. Interspecies scaling of dosages can be performed in a manner known in the art, e.g., as disclosed in Mordenti et al., Pharmaceut. Res., 8:1351 (1991).

When *in vivo* administration of an antagonist thereof is employed, normal dosage amounts may vary from about 10 ng/kg to up to 100 mg/kg of mammal body weight or more per day, preferably about 1 µg/kg/day to 10 mg/kg/day, depending upon the route of administration. Guidance as to particular dosages and methods of delivery is provided in the literature; see, for example, U.S. Pat. Nos. 4,657,760; 5,206,344; or 5,225,212. It is anticipated that different formulations will be effective for different treatment compounds and different disorders, that administration targeting one organ or tissue, for example, may necessitate delivery in a manner different from that to another organ or tissue. Those skilled in the art will understand that the dosage of antagonist that must be administered will vary depending on, for example, the mammal which will receive the agonist or antagonist, the route of administration, and other drugs or therapies being administered to the mammal.

Depending on the type of cells and/or severity of the disease, about 1 µg/kg to 15 mg/kg (e.g. 0.1-20 mg/kg) of antagonist antibody is an initial candidate dosage for administration, whether, for example, by one or more separate administrations, or by continuous infusion. A typical daily dosage might range from about 1 µg/kg to 100 mg/kg or more, depending on the factors mentioned above. For repeated administrations over several days or longer, depending on the condition, the treatment is sustained until a desired suppression of disease symptoms occurs. However, other dosage regimens may be useful.

A single type of antagonist may be used in the methods of the invention. For example, a WISP-1 immunoadhesin molecule may be administered. Alternatively, the skilled practitioner may opt to employ a combination of antagonists in the methods, e.g., a combination of a WISP-1 immunoadhesin and WISP-1 antibody. It is contemplated that yet additional therapies may be employed in the methods. The one or more other therapies may include but are not limited to, administration of radiation therapy, cytokine(s), growth inhibitory agent(s), chemotherapeutic agent(s), cytotoxic agent(s), tyrosine kinase inhibitors, ras farnesyl transferase inhibitors, angiogenesis inhibitors, and cyclin-dependent kinase inhibitors which are known in the art and defined further with particularity in Section I above. In addition, therapies based on therapeutic antibodies that target tumor antigens such as Rituxan™ or Herceptin™ as well as anti-angiogenic antibodies such as anti-VEGF.

Preparation and dosing schedules for chemotherapeutic agents may be used according to manufacturers' instructions or as determined empirically by the skilled practitioner. Preparation and dosing schedules for such chemotherapy are also described in *Chemotherapy Service* Ed., M.C. Perry, Williams & Wilkins, Baltimore, MD (1992). The chemotherapeutic agent may precede, or follow administration of, e.g. an antagonist, or may be given simultaneously therewith. The antagonist, for instance, may also be combined with an anti-oestrogen compound such as tamoxifen or an anti-progesterone such as onapristone (see, EP 616812) in dosages known for such molecules.

It may be desirable to also administer antibodies against other antigens, such as antibodies which bind to CD20, CD11a, CD18, CD40, CD44, ErbB2, EGFR, ErbB3, ErbB4, vascular endothelial factor (VEGF), or other TNFR family members (such as DR4, DR5, OPG, TNFR1, TNFR2). Alternatively, or in addition, two or more antibodies binding the same or two or more different antigens disclosed herein may be co-administered to the patient. Sometimes, it may be beneficial to also administer one or more cytokines to the patient. In one embodiment, the antagonists herein are co-administered with a growth inhibitory agent. For example, the growth

inhibitory agent may be administered first, followed by an antagonist of the present invention.

The antagonist and one or more other therapies may be administered concurrently or sequentially. Following administration of antagonist, treated cells in vitro can be analyzed. Where there has been in vivo treatment, a treated mammal can be monitored in various ways well known to the skilled practitioner.

ARTICLES OF MANUFACTURE

In another embodiment of the invention, an article of manufacture containing materials useful for the treatment of the disorders described above, is provided. The article of manufacture comprises a container and a label. Suitable containers include, for example, bottles, vials, syringes, and test tubes. The containers may be formed from a variety of materials such as glass or plastic. The container holds a composition which is effective for treating the condition and may have a sterile access port (for example the container may be an intravenous solution bag or a vial having a stopper pierceable by a hypodermic injection needle). The active agents in the composition may comprise antagonist(s). The label on, or associated with, the container indicates that the composition is used for treating the condition of choice. The article of manufacture may further comprise a second container comprising a pharmaceutically-acceptable buffer, such as phosphate-buffered saline, Ringer's solution and dextrose solution. It may further include other materials desirable from a commercial and user standpoint, including other buffers, diluents, filters, needles, syringes, and package inserts with instructions for use.

The following examples are offered for illustrative purposes only, and are not intended to limit the scope of the present invention in any way.

All patent and literature references cited in the present specification are hereby incorporated by reference in their entirety.

EXAMPLES

Commercially available reagents referred to in the examples were used according to manufacturer's instructions unless otherwise indicated. The source of those cells identified in the following examples, and throughout the specification, by ATCC accession numbers is the American Type Culture Collection, Manassas, VA. Unless otherwise noted, the present invention uses standard procedures of recombinant DNA technology, such as those described hereinabove and in the following textbooks: Sambrook et al., *Molecular Cloning: A Laboratory Manual*, Cold Spring Harbor Press N.Y., 1989; Ausubel et al., *Current Protocols in Molecular Biology*, Green Publishing Associates and Wiley Interscience, N.Y., 1989; Innis et al., *PCR*

Protocols: A Guide to Methods and Applications, Academic Press, Inc., N.Y., 1990; Harlow et al., *Antibodies: A Laboratory Manual*, Cold Spring Harbor Press, Cold Spring Harbor, 1988; Gait, M.J., *Oligonucleotide Synthesis*, IRL Press, Oxford, 1984; R.I. Freshney, *Animal Cell Culture*, 1987; Coligan et al., *Current Protocols in Immunology*, 1991.

A description of the assays and materials referred to in the following Examples is provided below:

Proteins and Antibodies: Full length human WISP-1-Fc fusion protein was expressed and purified as described previously (Desnoyers et al., *J. Biol. Chem.*, 276:47599-47607 (2001)). Purified mouse anti-rat CD44 antibody (clone OX49) was from BD Bioscience (Bedford MA). The mouse IgG2a isotype control antibody was from Pharmingen. The FITC conjugated mouse anti-rat CD44 was from Serotec (Raleigh, NC) and the FITC conjugated mouse IgG1 isotype control antibody was from DAKO (Denmark). The rat anti-mouse CD44 (clone KM114) was from Pharmingen. The actin antibody (clone C4) was from ICN biomedicals (Aurora, OH). Murine anti-human WISP-1 monoclonal antibodies were generated and selected using WISP-1-Fc as an immunogen and the protocols and reagents described in the Examples of US Patent No. 6,387,657 issued May 14, 2002. Five such monoclonal antibodies have been deposited by Applicants with the ATCC, as noted below.

Cells and Tissue Specimens: NRK-49F and SW480 cell lines (ATCC, Manassas, VA) were maintained in High glucose Dulbecco's Modified Eagle's Medium (HGMEM) supplemented with 10% fetal calf serum (FBS). Parental C57MG, C57MG expressing Wnt-1 (C57MG/Wnt-1), NRK-49F expressing high levels of hWISP-1 (NRK/WISP-1H), NRK-49F expressing lower levels of hWISP-1 (NRK/WISP-1L) and NRK-49F containing an empty vector (NRK/control) were obtained from Arnold Levine (Princeton University, Princeton, NJ) and maintained in media containing 2.5 µg ml⁻¹ puromycin (Xu et al., *Genes Dev.*, 14:585-595 (2000)). Wnt-1 transgenic mice obtained from Harold Varmus (National Cancer Institute, National Institutes of Health, Bethesda, MD) were bred with p53 null mice obtained from Jackson Laboratories (Bar Harbor, MA) to generate the Wnt-1 transgenic/p53 knock out mice. Specimens of breast tumor and mammary duct epithelium were excised from those mice for analysis.

Particle Exclusion Assay: The erythrocyte sedimentation assay was used as described previously, with minor modifications. (Knudson et al., *J. Cell Sci.*, 99:227-235 (1991)). NRK/WISP-1 (normal rat kidney fibroblasts transfected with WISP-1; see Xu et al., *Genes Dev.*, 14:585-595 (2000)) and NRK/control cells (normal rat kidney fibroblasts; obtained from Arnold Levine, Princeton University, Princeton, NJ) (1 X 10⁵ cells/well) were seeded in Falcon 6 well plates (Becton Dickinson, Franklin Lanes, NJ) and maintained in high glucose Dulbecco's modified Eagle's Medium supplemented

with 10% fetal bovine serum (FBS) (Life Technologies, Gaithersburg, MD) overnight at 37° C, 5% CO₂. The next day, the media was removed and 10⁸ sheep RBCs (Inter-Cell Technologies, Hopewell, NJ) in 750µl PBS/0.1% BSA (Boehringer Mannheim, Indianapolis, IN) were added and allowed to settle for 15 minutes. The cells were observed using a Nikon Diaphot 300 inverted microscope, and digital images were acquired using a SONY DKC-5000 digital photo camera (Japan).

Hyaluronan Staining of Cultured Cells: Hyaluronan

Staining was performed as previously described, with minor modifications (Pienimäki et al., *J. Biol. Chem.*, 276:20428-20435 (2001)). NRK/WISP-1 and NRK/control cells (8-10 X 10³) were plated in 8 well plastic chamber slides and maintained (as described above) overnight at 37°C, 5% CO₂. The next day the cells were washed with phosphate buffered saline (PBS) and fixed with 4% paraformaldehyde in PBS for 20 minutes at room temperature. The cells were washed with PBS and the non-specific binding sites were saturated for 30 minutes with distilled water containing 1% bovine serum albumin (BSA) and 0.1% Triton X-100 (Calbiochem, LaJolla, CA). Biotinylated HA binding protein (HA-BP) (Seikagaku America, Falmouth, MA) (5µg/ml in PBS/3% BSA) was added and incubated overnight at room temperature. The next day the cells were washed with PBS and incubated for 30 minutes with 1:1000 FITC-conjugated streptavidin (DAKO, Carpinteria, CA) in TBS and 1% BSA. The cells were washed in PBS, mounted using Vectashield (Burlingame, CA) containing 1 µg/ml Hoechst 33342 (Molecular Probes, Eugene, OR) and visualized under a Nikon Eclipse-800 fluorescent microscope. Images were acquired with a Photometrics 300 CCD camera operated at -40°C interfaced with an Apple G3 computer. The same procedure was also used to stain frozen sections of mammary tumors from Wnt-1 transgenic mice).

Real time RT-PCR: Relative RNA expression level was determined using real-time RT-PCR as previously reported (Winer et al., *Anal. Biochem.*, 270:41-49 (1999)). Total RNA was extracted from cells with Tri-Reagent using the manufacturer's protocol (Molecular Research Center). Specific primers and fluorogenic probes were designed and used to amplify and quantitate gene expression level. The genes specific signals were normalized to that of the glyceraldehyde-3-phosphate dehydrogenase housekeeping gene. Triplicate sets of data were averaged for each condition. All TaqMan RT-PCR reagents were purchased from Applied Biosystems (Foster City, CA). To test the effect of recombinant WISP-1 on gene expression, petri dishes (35 mm) were coated overnight at 4°C with 75 µg ml⁻¹ hWISP-1-Fc in PBS. The next day, cells (3 X 10⁵ cells/dish) were plated on the coated surface and the expression analysis was performed

and the plates were incubated overnight at 37°C. The next day the upper chamber was wiped with a cotton swab and the cells that migrated to the lower side of the insert were stained with the Diff-Quik Stain Kit (Dade Behring Inc.) and counted under the microscope. Triplicate sets of data were averaged for each condition.

In certain cases, the coated inserts were washed and treated for an additional 2 hours at room temperature.

In situ Hybridization: PCR primers were designed to amplify either a 740 bp fragment of murine WISP-1 spanning from nt 440-1180 of NM_018865 (upper- 5' GGCTGCCATCTGTGACCCA (SEQ ID NO:12) and lower- 5' CATAGGACCT GCCGGGAGAA A (SEQ ID NO:13)) or a 706 bp fragment of murine WISP-1 spanning from nt 204-910 of NM_018865 (upper- 5' GCCGTGGCAGTCCTGAGGG (SEQ ID NO:14) and lower- 5' CAGCACCGGG CATTGACGTT A (SEQ ID NO:15)) or a 464 bp fragment of murine CD44 spanning from nt 144-608 of M27129 (upper- 5' TGGAGAAAAATGGCCGCTAC A (SEQ ID NO:16)) and lower-5' TGGGGTGCTC TTCTCGATGG (SEQ ID NO:17)) or a 630 bp fragment of murine HAS2 spanning from nt 927-1557 of NM_008216 (upper- 5' GGACAAATCGGCCACGTACA T (SEQ ID NO:18) and lower-5' CTTGCTCCAT CGGGTCTGC (SEQ ID NO:19)) . Primers included extensions encoding 27-nucleotide T7 or T3 RNA polymerase initiation sites to allow in vitro transcription of sense or antisense probes, respectively, from the amplified products. (Lu et al., Cell Vision, 1:169-176 (1994)). All tissues were fixed in 4% formalin and paraffin-embedded. Sections 3-5 microns thick were deparaffinized, deproteinized in 4 mg/ml of proteinase K for 30 minutes at 37 °C, and further processed for in situ hybridization as previously described. (Holcomb et al., EMBO J., Aug 2000 1;19(15):4046-55). ³³P-UTP labeled sense and antisense probes were hybridized to the sections at 55°C overnight. Unhybridized probe was removed by incubation in 20mg/ml RNase A for 30 minutes at 37°C, followed by a high stringency wash at 55 °C in 0.1 X SSC for 2 hours and dehydration through graded ethanols. The slides were dipped in NBT2 nuclear track emulsion, exposed in sealed plastic slide boxes containing dessicant for 4 weeks at 4 °C, developed and counterstained with hematoxylin and eosin.

EXAMPLE 1

Stable fibroblast cell lines expressing high (NRK/WISP-1H) or low (NRK/WISP-1L) levels of WISP-1 and a control cell line containing an empty vector (NRK/control) were used to evaluate the effect of WISP-1 on HA production. As demonstrated by the particle exclusion assay, NRK/WISP-1H cells accumulated a large hyaluronan pericellular coat capable of excluding sedimenting erythrocytes (Fig. 1a). NRK/WISP-1L accumulated a smaller pericellular matrix (Fig. 1b) whereas no matrix surrounded NRK/control (Fig. 1c) or NRK/WISP-1H cells treated with hyaluronidase (data not shown).

Accumulation of cell surface associated HA was also evaluated by fluorescent staining using the biotinylated HA binding protein (bHABP). The staining revealed HA accumulation at the NRK/WISP-1H cell surface (Fig. 1d) whereas no staining was detected on NRK control cells (Fig. 1e).

5 The effect of WISP-1 on HA secretion was evaluated by comparing the accumulation of HA in the culture media of NRK/WISP-1H to NRK/control cells over time. After 24 hours, HA concentration in the NRK/WISP-1H media was 3.5 fold greater in NRK/control cell media and gradually increased to 8 fold after 144 h (fig. 1f). Together these results show that WISP-1
10 expression in NRK cells promotes HA accumulation at the cell surface and in the culture media.

EXAMPLE 2

To identify the enzyme responsible for the WISP-1 triggered HA
15 production, the expression of the known HA synthases (HAS1, HAS2 and HAS3) in NRK/WISP-1H, NRK/WISP-1L and NRK/control cells was analyzed. HAS2 expression was increased up to 10 fold in WISP-1 producing cells whereas HAS1 and HAS3 mRNA levels were identical to the control (Fig. 2a). CD44 and RHAMM mRNA levels were also increased in NRK/WISP-1 cell lines whereas
20 hyaluronidase expression remained unchanged (Fig 2a). Moreover, HAS2, CD44 and RHAMM mRNA levels were proportional to WISP-1 expression. As demonstrated by FACS analysis (Fig. 2b) and Western blot (Fig. 2c), the increase in CD44 mRNA expression in the NRK/WISP-1H cells resulted in a 4- fold increase in CD44 protein level.

25

EXAMPLE 3

HA was shown to stimulate cell migration by interacting with two cell surface receptors, CD44 and RHAMM (Hall et al., supra; Bourguignon et al., supra). Because WISP-1 increased both HA production and CD44 and RHAMM
30 expression, the motility of WISP-1 expressing cells was evaluated. When plated at a low density, NRK/control cells proliferated and formed well defined colonies (Fig 3a). Proliferating NRK/WISP-1L cells formed less defined groups as some cells departed from the growing colonies (Fig 3b). No NRK/WISP-1H colonies were seen, cells scattered in a random pattern (Fig
35 3c). Contrary to the control (Fig. 3d), WISP-1 expressing cells also revealed a hyper elongated morphology with extended lamellipodia characteristic of high motility (Fig. 3e). Using time lapse microscopy, a 4 fold increase in the migration distance of NRK/WISP-1H was observed compared to control cells (Fig. 3f). NRK/WISP-1H also showed a 2.5 fold
40 increase in migration area in a cell wound healing assay (Fig. 3g). Together these results demonstrate that WISP-1 expression may promote cell migration.

EXAMPLE 4

To determine whether the ectopic addition of WISP-1 could promote HAS2 expression and cell migration, an assay was conducted using purified recombinant WISP-1-IgG (Desnoyers et al., supra). WISP-1 addition to the culture media did not promote HAS2 expression (data not shown). When NRK cells were plated onto a WISP-1-IgG coated surface, HAS2 expression increased compared to cells plated onto uncoated plastic or onto an irrelevant IgG chimeric protein (TNFR-IgG) coated surface (Fig. 4a). Although decorin alone promoted HAS2 expression, this induction was further increased three fold after the coated surface was incubated with WISP-1. The effects of ectopic WISP-1 addition on NRK cell migration were also examined. In a transwell assay, WISP-1-IgG induced the haptotactic migration of NRK cells when coated on the filter's lower surface (Fig. 4b). Coating of an irrelevant IgG chimeric protein (TNFR-IgG) or addition of WISP-1-IgG to the lower chamber did not promote migration. A CD44 or a WISP-1 antibody (Fig. 4b) inhibited the haptotactic migration induced by WISP-1. Moreover, WISP-1 haptotactic activity was not limited to fibroblasts as it also promoted the migration of SW480 cells, a colon adenocarcinoma cell line (Fig. 4c). Together these results demonstrate that ectopic WISP-1 addition increases HAS2 expression and promotes cell migration through a CD44 mediated mechanism. It is believed, though not fully understood, that WISP-1 presentation may be important for migration as it may need to be tethered to a substrate to elicit this activity.

EXAMPLE 5

Because WISP-1 is believed to be a downstream effector of Wnt-1, WISP-1, HAS2 and CD44 expression in a mammary epithelial cell line stably transfected with Wnt-1 (C57MG/Wnt-1) were analyzed. When compared to the control cell line (C57MG), C57MG/Wnt-1 cells showed a 2.7, 5.8 and 3 fold increase in WISP-1, HAS2 and CD44 mRNA expression respectively (Fig. 5a). As demonstrated by Western blot, C57/Wnt-1 cells also showed a 6-fold increase in CD44 protein content (Fig. 5b). Moreover, unlike the control cell line, Wnt-1 expressing cells failed to form distinct colonies and scattered when put in culture at low density (Fig. 5c). These results show that HAS2 and CD44 expression and cell motility are elevated in cell lines where WISP-1 expression is triggered by Wnt-1.

EXAMPLE 6

Wnt-1 expression in mammary epithelium promotes tumor development in transgenic mice (Li et al., Oncogene, 19:1002-1009 (2000); Tsukamoto et al., Cell, 55:619-625. (1988)). Because WISP-1 is a putative downstream

effector of Wnt-1, the mRNA expression of HAS2 and CD44 in spontaneous mammary tumors from MMTV-Wnt-1 transgenic mice was measured. When compared to mammary duct epithelium, HAS2 mRNA expression was increased between 2.5 and 5.5 fold in all mammary tumors analyzed (n=5; Fig. 5d). Similarly, CD44 mRNA expression was induced between 2.2 and 4.2 fold in all tumors (n=5; Fig 5e). These results show that HAS2 and CD44 are overexpressed in MMTV-Wnt-1 transgenic mice mammary tumors expressing WISP-1.

EXAMPLE 7

In situ hybridization demonstrated elevated WISP-1 expression in the peritumoral stroma of MMTV-Wnt-1 transgenic mouse mammary tumors (Fig. 6a, b). In contrast, HAS2 (Fig. 6c-d) and CD44 (Fig. 6 e-f) expression was found only in tumoral epithelial cells. The localization of CD44 in the tumor parenchyma was confirmed by immunohistochemistry (Fig. 6g). Link protein staining revealed an accumulation of hyaluronan associated with the tumor stroma (Fig. 6h). The highest staining intensity was found in the immediate vicinity of the tumor lobule whereas weaker staining was localized to the normal mammary duct epithelium. These findings are consistent with the suggestion that WISP-1 may regulate interactions between tumor and stromal cells that involve hyaluronic acid and CD44.

EXAMPLE 8

The metastasis and growth potential of transformed lines was evaluated as described previously (Welch et al., Cancer Res., 60:1552-1556 (2000)). Nine-week-old Swiss nu/nu female mice were used. Briefly, cells were harvested by trypsinization and washed twice with PBS. Each mouse was injected into the lateral tail vein with 100 μ l of a suspension containing 5×10^4 or 2.5×10^5 cells. At 2, 3 and 4 week postinjection, mice were examined using cine-magnetic resonance imaging (MRI) for apparition of pulmonary lesions and necropsies were performed. The lungs were perfused in Bouin's fixative, excised and H&E stained sections were generated for evaluation of pulmonary tumor colonization.

Longitudinal sections of the left lung, and a single transverse section of the cranial, medial, caudal, and accessory lobes of the right lung were evaluated. Histologically, affected lungs had spindloid neoplastic cells that infiltrated the pulmonary interstitium. However, because the severity of the changes was variable the following grading system was established.

I = Minimal involvement of the pulmonary interstitium - 10-20% of the lung is affected. Nests and clusters of spindloid neoplastic cells multifocally disrupt the pulmonary interstitium. Affected foci are commonly along the

pleural surface, extending into the interstitium. No vessels or large airways are affected.

II = Moderate involvement of the pulmonary interstitium - 20 - 50% of the lung is affected. Nests and clusters of spindloid neoplastic cells

5 multifocally disrupt the pulmonary interstitium. Some vessels or large airways are filled with spindloid cells. Spindloid cells frequently form a broad band or large mass subjacent to the pleura and extending into the interstitium. Overlying mesothelium is plump (reactive).

10 III = Severe involvement of the pulmonary interstitium - 50- 100% of the lung is affected. Nests and clusters of spindloid neoplastic cells multifocally disrupt the pulmonary interstitium. Blood vessels and large airways are filled with spindloid cells. Spindloid cells form a broad band or large mass subjacent to the pleura and extending into the interstitium. Spindloid cells are often embedded in a pale basophilic to amphophilic

15 acellular material. Overlying mesothelium is plump (reactive). Minimal unaffected area for gas exchange remains.

The results of the lung colonization of nude mice inoculated with NRK/WISP-1H, NRK/WISP-1L and NRK/control cells are summarized in Figures 7 and 10. In mice inoculated with NRK/WISP-1H or NRK/WISP-1L, cells formed

20 lung masses in a time- and dose-dependent manner. The most severe lesions were seen in mice injected with 2.5×10^5 NRK/WISP-1H and necropsied at 4 weeks post-injection. At 2 and 3 weeks post-injection, the mice had less severe but progressive tumor formation (grade I - II; see grading system above). Mice injected with 2.5×10^5 NRK/WISP-1L had only minimal lung

25 tumor formation by 4 weeks post-injection. After 4 weeks post-injection, mice inoculated with 0.5×10^5 NRK/WISP-1L cells were either normal or had minimal neoplastic infiltration (grade I), whereas mice injected with 0.5×10^5 NRK/WISP-1H cells showed increased neoplastic infiltration (grade I-II). Histological observation revealed that the lungs of animals

30 inoculated with NRK/control cells were normal (Fig. 7a-c). In NRK/WISP-1 injected animals, neoplastic spindloid cells initially formed small clusters within, protruding from, or subjacent to the pleural surface of the lungs (grade I; Fig. 7d-f) often embedded in a pale basophilic to amphophilic acellular material. In more severely affected animals, the

35 neoplastic cells formed a broad confluent band subjacent to the pleura (grade II; Fig. 7g-i). In the most severely affected animals, blood vessels and large airways were filled with spindloid cells. (grade III; Fig. 7j-l). MRI analysis showed increased density (hypersignal) outlining the right and left lungs along the pleura in animals with histologic grade

40 II and grade III invasion. Severity of the lesion on MRI was consistent with the histologic score (Figure 7a, d, g, j). These results indicate that WISP-1 expression promotes cellular metastatic growth potential.

በጥንቃቄ ሲገባ ለጥያቄው ምክር ቤቱ ማቅረብ ይገባል፡፡

5

10

15

20

3

4

lysate. The WISP-1 antibodies 6F8, 3A7, 10H12, 3A11, 6E3, 3H10, 5G1, and 10B1 recognized both human and murine WISP-1 only when generated with the baculovirus expression system (Fig. 11b). These antibodies did not recognize human WISP-1 when produced in a bacterial or mammalian expression system. The antibody from clone 9C10 did not bind to any protein after Western blot.

These results suggest that WISP-1 antibodies 3D11, 11C2, 9C11 and 5D4 specifically recognize human WISP-1 and can be used for WISP-1 detection by Western blot.

EXAMPLE 11

An assay was conducted to identify the epitopes recognized by the WISP-1 antibodies 11C2, 9C11, 5D4 and 3D11.

Full length human WISP-1 (GenBank accession number AF100779) was cloned into a pIRESpuro2 expression vector (Clontech Laboratories, Palo Alto, CA) encoding 6 histidines downstream of the WISP-1 sequence. Deletion mutants were also generated by removing one, two or three domains of human WISP-1. The resulting constructs were also cloned into the pIRESpuro2 expression vector. The nomenclature used to identify the different WISP-1 constructs refer to the domains they contain. (see Fig. 12B) Domain 1 is the insulin-like growth factor binding protein domain (IFGBP), domain 2 is the von Willebrand factor C (VWFc) domain, domain 3 is the thrombospondin (TSP) domain, and the domain 4 is the C-terminal (CT) domain. The variable region resides between domain 2 and 3. These regions and domains of WISP-1 are illustrated in Fig. 12A.

The sequences encoding these domains of WISP-1 are as follows:

Sequences of WISP-1 Constructs

Domain 1:

GAATTCACCATGAGGTGGTTCCTGCCCTGGACGCTGGCAGCAGTGACAGCAGCAGCCGCCAGCACCGTCCTGGCC
ACGGCCCTCTCTCCAGCCCCTACGACCATGGACTTTACTCCAGCTCCACTGGAGGACACCTCCTCACGCCCCCA
TTCTGCAAGTGGCCATGTGAGTGCCCGCCATCCCCACCCGCTGCCCCGCTGGGGGTGAGCCTCATCACAGATGGC
TGTGAGTGCTGTAAGATGTGCGCTCAGCAGCTTGGGGACAACCTGCACGGAGGCTGCCATCTGTGACCCCCACCGG
GGCCTCTACTGTGACTACAGCGGGGACCGCCCCGAGGTACGCAATAGGAGTGTGTGCACAGGCGGCCGACACCAC
CATCACCATCACCATCACTAAGTGAGGCCGATAGATAACTGATCCAGTGTGCTGGAATTAATTC (SEQ ID
NO:3)

Domain 2:

GAATTCACCATGAGGTGGTTCCTGCCCTGGACGCTGGCAGCAGTGACAGCAGCAGCCGCCAGCACCGTCCTGGCC
ACTGCAGTGGTGGTGTGGGCTGCGTCTGGATGGGGTGGCTACAACAACGGCCAGTCCCTCCAGCCTAACTGC
AAGTACAACCTGCACGTGCATCGACGGCGGGTGGGCTGCACACCACTGTGCCTCCGAGTGGCCCCCGGCTCTC
TGGTGCCCCCACCGCGCGGTGAGCATACTGGCCACTGCTGTGAGCAGTGGGTATGTGCGGCCGCGACACCAC
CATCACCATCACCATCACTAAGTGAGGCCGATAGATAAC (SEQ ID NO:4)

Domain 3:

GAATTCACCATGAGGTGGTTTCCTGCCCTGGACGCTGGCAGCAGTGACAGCAGCAGCCGCCAGCACCGTCCTGGCC
 ACTGCAGCATGGCACAGGAAGTGCATAGCCTACACAAGCCCCCTGGAGCCCTTGCTCCACCAGCTGCGGCCCTGGGG
 5 GTCTCCACTCGGATCTCCAATGTTAACGCCCAGTGCTGGCCTGAGCAAGAGAGCCGCCTCTGCAACTTGCGGCCA
 TGCGATGTGGACATCCATACACTCATTAAAGGCGGCCGCACACCACCATCACCATCACCATCACTAAGTGAGGCCG
 CATAGATAACTGATCCAGTGT (SEQ ID NO:5)

Domain 4:

10 GAATTCACCATGAGGTGGTTTCCTGCCCTGGACGCTGGCAGCAGTGACAGCAGCAGCCGCCAGCACCGTCCTGGCC
 ACTGCAGGGAAGAAGTGTCTGGCTGTGTACCAGCCAGAGGCATCCATGAACTTCACACTTGCGGGCTGCATCAGC
 ACACGCTCCTATCAACCCAAGTACTGTGGAGTTTGCATGGACAATAGGTGCTGCATCCCCTACAAGTCTAAGACT
 ATCGACGTGTCCTTCCAGTGTCTGTGGGCTTGGCTTCTCCCGCCAGGTCCTATGGATTAATGCCTGCTTCTGT
 AACCTGAGCTGTAGGAATCCCAATGACATCTTTGCTGACTTGGAATCCTACCCTGACTTCTCAGAAATTGCCAAC
 15 GCGGCCGCACACCACCATCACCATCACCATCACTAAGTGAGGCCGCATAGATAACTGATCCAGTGTG (SEQ
 ID NO:6)

Domain 1,2:

GAATTCACCATGAGGTGGTTTCCTGCCCTGGACGCTGGCAGCAGTGACAGCAGCAGCCGCCAGCACCGTCCTGGCC
 20 ACGGCCCTCTCTCCAGCCCCCTACGACCATGGACTTTACTCCAGCTCCACTGGAGGACACCTCCTCACGCCCCCAA
 TTCTGCAAGTGGCCATGTGAGTGCCCGCCATCCCCACCCCGCTGCCCGCTGGGGGTGAGCCTCATCACAGATGGC
 TGTGAGTGCTGTAAGATGTGCGCTCAGCAGCTTGGGGACAACGACGAGGCTGCCATCTGTGACCCCCACCGG
 GGCCTCTACTGTGACTACAGCGGGGACCGCCCCGAGGTACGCAATAGGAGTGTGTGCACAGGTGGTCGGTGTGGGC
 TGCGTCCTGGATGGGGTGCCTTACAACAACGGCCAGTCCTTCCAGCCTAACTGCAAGTACAACGACGTGCATC
 25 GACGGCGCGGTGGGCTGCACACCACTGTGCCTCCGAGTGCGCCCCCGCGTCTCTGGTGCCCCCACCGCGGCGC
 GTGAGCATACCTGGCCACTGCTGTGAGCAGTGGGTATGTGCGGCCGCACACCACCATCACCATCACCATCACTAA
 GTGAGGCCGCATAGATAAC (SEQ ID NO:7)

Domain 1,2,3:

30 GAATTCACCATGAGGTGGTTTCCTGCCCTGGACGCTGGCAGCAGTGACAGCAGCAGCCGCCAGCACCGTCCTGGCC
 ACGGCCCTCTCTCCAGCCCCCTACGACCATGGACTTTACTCCAGCTCCACTGGAGGACACCTCCTCACGCCCCCAA
 TTCTGCAAGTGGCCATGTGAGTGCCCGCCATCCCCACCCCGCTGCCCGCTGGGGGTGAGCCTCATCACAGATGGC
 TGTGAGTGCTGTAAGATGTGCGCTCAGCAGCTTGGGGACAACGACGAGGCTGCCATCTGTGACCCCCACCGG
 GGCCTCTACTGTGACTACAGCGGGGACCGCCCCGAGGTACGCAATAGGAGTGTGTGCACAGGTGGTCGGTGTGGGC
 35 TGCGTCCTGGATGGGGTGCCTTACAACAACGGCCAGTCCTTCCAGCCTAACTGCAAGTACAACGACGTGCATC
 GACGGCGCGGTGGGCTGCACACCACTGTGCCTCCGAGTGCGCCCCCGCGTCTCTGGTGCCCCCACCGCGGCGC
 GTGAGCATACCTGGCCACTGCTGTGAGCAGTGGGTATGTGAGGACGACGCCAAGAGGCCACGCAAGACCGCACCC
 CGTGACACAGGAGCCTTCGATGCTGTGGGTGAGGTGGAGGCATGGCACAGGAAGTGCATAGCCTACACAAGCCCC
 TGGAGCCCTTGCTCCACCAGCTGCGGCCCTGGGGGTCTCCACTCGGATCTCCAATGTTAACGCCCAGTGCTGGCCT
 40 GAGCAAGAGAGCCGCCTCTGCAACTTGCGGCCATGCGATGTGGACATCCATACACTCATTAAAGGCGGCCGCACAC
 CACCATCACCATCACCATCACTAAGTGAGGCCGCATAGATAACTGATCCAGTGTGCTGGA (SEQ ID NO:8)

Domain 1,2,4:

GAATTCACCATGAGGTGGTTTCCTGCCCTGGACGCTGGCAGCAGTGACAGCAGCAGCCGCCAGCACCGTCCTGGCC
ACGGCCCTCTCTCCAGCCCCCTACGACCATGGACTTTACTCCAGCTCCACTGGAGGACACCTCCTCACGCCCCCAA
TTCTGCAAGTGGCCATGTGAGTGCCCGCCATCCCCACCCCGCTGCCCCTGGGGGTGAGCCTCATCACAGATGGC
5 TGTGAGTGCTGTAAGATGTGCGCTCAGCAGCTTGGGGACAACCTGCACGGAGGCTGCCATCTGTGACCCACCAGG
GGCCTCTACTGTGACTACAGCGGGGACCGCCGAGGTACGCAATAGGAGTGTGTGCACAGGTGGTGGTGTGGGC
TGCGTCTGGATGGGGTGGCTACAACAACGGCCAGTCCTTCCAGCCTAACTGCAAGTACAACCTGCACGTGCATC
GACGGCGCGGTGGGTGCACACCACTGTGCCTCCGAGTGCGCCCCCGCTCTTGGTGGCCCCACCCGCGGCGC
GTGAGCATACCTGGCCACTGCTGTGAGCAGTGGGTATGTCTGCAGGCAGGGAAGAAGTGTCTGGCTGTGTACCAG
10 CCAGAGGCATCCATGAACCTTACACTTGCAGGCTGCATCAGCACACGCTCCTATCAACCCAAGTACTGTGGAGTT
TGCATGGACAATAGGTGCTGCATCCCCCTACAAGTCTAAGACTATCGACGTGTCTTCCAGTGTCTGATGGGCTT
GGCTTCTCCCGCCAGGTCTTATGGATTAATGCCTGCTTCTGTAACCTGAGCTGTAGGAATCCCAATGACATCTTT
GCTGACTTGAATCCTACCTGACTTCTCAGAAATTGCCAACGCGGCCGCACACCACCATCACCATCACCATCAC
TAAGTGAGGCCGCATAGATAACTGATCCAGTGTGCTGGAATTAATTGCTGTCTGCGAGGGCCAGCTGTTGGGGT
15 GAGTACTCCCTCTCAAAGCGGGCATGACTTCTGCGCTA (SEQ ID NO:9)

Domain 1,3,4:

GAATTCACCATGAGGTGGTTTCCTGCCCTGGACGCTGGCAGCAGTGACAGCAGCAGCCGCCAGCACCGTCCTGGCC
ACGGCCCTCTCTCCAGCCCCCTACGACCATGGACTTTACTCCAGCTCCACTGGAGGACACCTCCTCACGCCCCCAA
20 TTCTGCAAGTGGCCATGTGAGTGCCCGCCATCCCCACCCCGCTGCCCCTGGGGGTGAGCCTCATCACAGATGGC
TGTGAGTGCTGTAAGATGTGCGCTCAGCAGCTTGGGGACAACCTGCACGGAGGCTGCCATCTGTGACCCACCAGG
GGCCTCTACTGTGACTACAGCGGGGACCGCCGAGGTACGCAATAGGAGTGTGTGCGCATGCTGTGGGTGAGGTG
GAGGCATGGCACAGGAACCTGCATAGCCTACACAAGCCCTGGAGCCCTTGCTCCACCAGCTGCGGCCCTGGGGGTG
TCCACTCGGATCTCCAATGTTAACGCCAGTGCTGGCCTGAGCAAGAGAGCCGCTCTGCAACTTGCAGGCCATGC
25 GATGTGGACATCCATACACTCATTAAAGGCAGGGAAGAAGTGTCTGGCTGTGTACCAGCCAGAGGCATCCATGAAC
TTCACACTTGCAGGCTGCATCAGCACACGCTCCTATCAACCCAAGTACTGTGGAGTTTGCATGGACAATAGGTGC
TGATCCCCCTACAAGTCTAAGACTATCGACGTGTCTTCCAGTGTCTGATGGGCTTGGCTTCTCCCGCCAGGTG
CTATGGATTAATGCCTGCTTCTGTAACCTGAGCTGTAGGAATCCCAATGACATCTTTGCTGACTTGAATCCTAC
CCTGACTTCTCAGAAATTGCCAACGCGGCCGCACACCACCATCACCATCACCATCACTAAGTGAGGCCGCATAGA
30 TAAC (SEQ ID NO:10)

Domain 2,3,4:

GAATTCACCATGAGGTGGTTTCCTGCCCTGGACGCTGGCAGCAGTGACAGCAGCAGCCGCCAGCACCGTC
CTGGCCACTGCAGTGGTGGTGTGGGCTGCGTCTGGATGGGGTGGCTACAACAACGGCCAGTCCTTCCAGCCT
35 AACTGCAAGTACAACCTGCACGTGCATCGACGGCGGCTGGGCTGCACACCACTGTGCCTCCGAGTGCGCCCCCG
CGTCTCTGGTGGCCCCACCCGCGCGGTGAGCATACCTGGCCACTGCTGTGAGCAGTGGGTATGTGAGGACGAC
GCCAAGAGGGCCACGCAAGAGCGCACCCCGTGACACAGGAGCCTTCGATGCTGTGGGTGAGGTGGAGGCATGGCAC
AGGAACTGCATAGCCTACACAAGCCCTGGAGCCCTTGCTCCACCAGCTGCGGCCCTGGGGGTCTCCACTCGGATC
TCCAATGTTAACGCCAGTGCTGGCCTGAGCAAGAGAGCCGCTCTGCAACTTGCAGGCCATGCGATGTGGACATC
40 CATACTCATTAAGGCAGGGAAGAAGTGTCTGGCTGTGTACCAGCCAGAGGCATCCATGAACCTTACACTTGGC
GGCTGCATCAGCACACGCTCCTATCAACCCAAGTACTGTGGAGTTTGCATGGACAATAGGTGCTGCATCCCCCTAC
AAGTCTAAGACTATCGACGTGTCTTCCAGTGTCTGATGGGCTTGGCTTCTCCCGCCAGGTCTATGGATTAAT

GCCTGCTTCTGTAACTGAGCTGTAGGAATCCCAATGACATCTTTGCTGACTTGGAAATCCTACCCTGACTTCTCA
GAAATTGCCAAACGCGCCGCACACCACCATCACCATCACCATCAGTGAAGTGAGGCCGCATAGATAACTGATCCAG
TGTGCTGGAATTAATTCGCTGTCTGCGA (SEQ ID NO:11)

5 Cells (HEK 293T) were transfected with the different constructs, and
the culture media was collected after 48 hours. One milliliter of culture
media was incubated with 20 µl of cobalt-agarose for 1 hour, centrifuged
and washed. The adsorbed proteins were eluted by heating the pellet at 100
°C for 5 minutes in 20 µl of SDS-PAGE sample buffer. The samples were
10 electrophoresed, electro-transferred onto PVDF and probed with the
different WISP-1 antibodies.

Antibodies 11C2, 9C11 and 5D4 recognized only WISP-1 constructs
containing the 19 first amino acids of the variable region located between
domain 2 and 3 (Fig. 12C; 12E; 12G). The WISP-1 antibody 3D11 recognized
15 only WISP-1 constructs containing the domain 1 (amino acids 24 to 117; Fig.
12D; 12F).

These results indicate that the antibodies 11C2, 9C11 and 5D4
recognize specifically the variable region of WISP-1 whereas the antibody
3D11 recognizes specifically the domain 1 of WISP-1.

20

EXAMPLE 12

An assay was conducted to identify the epitope recognized by the
WISP-1 antibody 9C10.F5 (also referred to herein as "9C10").

Culture media from HEK 293T cells transfected with the various WISP-1
25 deletion constructs (as described above in Example 11) was incubated with 1
µg of WISP-1 antibody 9C10 and 20 µl of protein A-agarose for 1 hour at
room temperature. The immunocomplex was precipitated by centrifugation and
eluted by heating the pellet at 100 °C for 5 minutes in 20 µl of SDS-PAGE
sample buffer. The samples were electrophoresed, electro-transferred onto
30 PVDF and probed with WISP-1 antibody 11C2.

The antibody 9C10 immunoprecipitated only constructs containing the
domain 1 of WISP-1 (Fig. 13). These results demonstrate that the antibody
9C10 specifically recognizes the domain 1 of WISP-1 and can be used for
immunoprecipitation.

35

EXAMPLE 13

WISP-1 antibody 9C10 (100 µl of 2 µg/ml in carbonate buffer, pH 9.6)
was coated to Maxisorb plates overnight at 4 °C. The plates were blocked
with 200 µl of PBS/3% BSA for 1 hour. A standard curve was made of serial
40 dilutions of WISP-1-Fc (100 µl in PBS/3% BSA) and incubated for 1 hour.
After the incubation, the plates were washed with 100 µl PBS/0.05% Tween
and WISP-1 antibodies (100 µl of 2 µg/ml) in PBS/3% BSA (biotinylated 11C2

5 of 55B) were incubated for 1 hour. For biotinylated 11C2, the plates were further incubated with 2 µg/ml HRP-conjugated streptavidin. For 55B, the plates are washed and incubated with HRP-conjugated donkey anti-rabbit IgG for 1 hour. At the end of the incubation, the wells were washed 6 times with 200 µl of PBS containing 0.05% Tween-20, and the signal was visualized using 100 µl of the horseradish peroxidase chromogenic substrate TMB (Kirkegaard & Perry Laboratories). The reaction was stopped with 100 µl of 1 M phosphoric acid, and the OD at 450 nm was measured. Non-specific binding was determined in parallel incubations by omitting microtiter well coating. No signal was generated when WISP-1-Fc or a WISP-1 antibody was omitted.

15 Using the antibody 9C10 for capture and the antibodies 11C2 and 55B for detection, an ELISA was conducted capable of detecting concentration of WISP-1 as low as 0.4 µg/ml (Fig. 14). This ELISA may be useful for detecting WISP-1 protein in biological fluids such as serum.

EXAMPLE 14

20 Maxisorb plates were coated overnight at 4 °C with 50 µl/well of 10 µg/ml heparin (Sigma). The non specific binding sites were blocked with 200µl of PBS/3% BSA for 1 hour. The plates were then incubated for 1 hour with 50 µl of 6 µg/ml hWISP-1-Fc in PBS/3% BSA in the presence of serial dilutions of WISP-1 antibodies. The plates were washed with PBS/0.05% Tween and further incubated 1 hour with 50 µl of 2 µg/ml HRP conjugated anti-human IgG-Fc in PBS/3% BSA. The plates were washed, and 100 µl of HRP substrate (TMB) was added. The color development was stopped with 100 µl of 1 M phosphoric acid and the OD at 450 nm was measured.

25 The WISP-1 antibodies 11C2, 5D4 and 9C11 inhibited WISP-1 binding to heparin with an IC₅₀ of 1.9, 2.5 and 3.7 µg/ml, respectively (Fig. 15). The antibody 3D11 moderately reduced WISP-1 binding to heparin with a maximal inhibition of 62% at the highest concentration tested (40 µg/ml). The antibody 9C10 did not attenuate WISP-1 heparin binding, showing an inhibition curve similar to the irrelevant antibody control.

35 These results demonstrate that antibodies recognizing the variable region can inhibit WISP-1 binding to heparin. Because the two WISP-1 antibodies recognizing domain 1 have little or no effect on WISP-1 binding to heparin, it is presently believed that the domain 1 is less likely to participate in this interaction.

EXAMPLE 15

40 Haptotaxis was measured using a modified Boyden chamber system. The underside of 8 µm porosity 24-well format PET membrane filters (Becton Dickinson, Franklin Lakes, New Jersey) were coated overnight at 4 °C with

50 μ l of protein (50 μ g/ml) in PBS. Normal rat kidney cells (NRK; 5×10^4 /0.5 ml HGD MEM/10% FBS) were added to the upper chamber, the lower chamber was filled with the same media and the plates were incubated at 37 °C. The next day, the upper chamber was wiped with a cotton swab, and the cells that migrated to the lower side of the insert were stained and counted under a microscope. Triplicate sets of data were averaged for each condition. In certain cases, the coated inserts were washed and treated for an additional 2 hours at room temperature.

In a transwell assay, WISP-1-Fc coated on the filters' lower surface induced the haptotactic migration of NRK cells (Fig. 16A). Coating of an irrelevant IgG chimeric protein (TNFR-Fc) or addition of WISP-1-Fc to the lower chamber did not promote migration. In the presence of coated WISP-1-Fc, five different WISP-1 antibodies (9C10, 11C2, 3D11, 9C11, 5D4) markedly inhibited cell migration. In the absence of coated WISP-1-Fc, these antibodies did not show any effect on cell migration. These results demonstrate that WISP-1 antibodies can modulate cell migration in the presence of WISP-1. By blocking cell migration, WISP-1 antibodies may play an important therapeutic role in preventing cancer progression.

A summary of the characteristics and properties of the 3D11, 9C10, 11C2, 5D4, and 9C11 antibodies discussed in the Examples above is provided in Fig. 16B.

EXAMPLE 16

Nine-week-old Swiss nu/nu female mice were injected into the lateral tail vein with 100 μ l of a suspension containing 2.5×10^5 cells. Starting on the day of the inoculation, the mice were treated semiweekly by intraperitoneal injection (10 mg/kg) of WISP-1 antibodies (9C11, 11C2, 5D4, 9C10, 3D11) or isotype control antibody. After 3 weeks, lungs were perfused with 4% neutral buffered formalin, excised and H&E stained sections were generated. Longitudinal sections of the left lung, and a transverse section of the cranial, medial, caudal, and accessory lobes of the right lung were evaluated. For each slide the number of metastatic foci was counted and the average area of metastatic foci was measured using the SPOT RT software (Diagnostic Instruments Incorporated, Burlington, California). The area (μ m²) was determined for at least five individual metastatic foci in four sections of lung.

After 3 weeks, the severity of the lesions found in WISP-1 antibody treated animals was greatly attenuated compared to control (Fig. 17 a, b). The number of nodules and the average area of the metastatic foci found in mice treated with WISP-1 antibodies (n=5) were reduced compared to animals treated with a control antibody (Fig. 17 c,d). Furthermore, upon WISP-1 antibody treatment, the total pulmonary area covered by the lesions was

reduced by 82-97% compared to animal treated with an isotype control antibody (Fig. 17e). These results demonstrate the *in vivo* efficacy of WISP-1 antibodies at reducing the tumor burden related to metastasis. Although the mechanism of action of the WISP-1 antibodies is not fully understood, it is believed their efficacy may be mediated by a growth reducing ability and/or a capacity at inhibiting the motility, invasion and seeding of cancer cells at a tissue site.

Deposit of Material

The following materials have been deposited with the American Type Culture Collection, 10801 University Blvd., Manassas, VA 20110-2209, USA (ATCC):

Material	ATCC Dep. No.	Deposit Date
3D11.D7	_____	Sept. __, 2002
11C2.C10	_____	Sept. __, 2002
9C10.F5	_____	Sept. __, 2002
5D4.F6	_____	Sept. __, 2002
9C11.C7	_____	Sept. __, 2002

This deposit was made under the provisions of the Budapest Treaty on the International Recognition of the Deposit of Microorganisms for the Purpose of Patent Procedure and the Regulations thereunder (Budapest Treaty). This assures maintenance of a viable culture of the deposit for 30 years from the date of deposit. The deposit will be made available by ATCC under the terms of the Budapest Treaty, and subject to an agreement between Genentech, Inc. and ATCC, which assures permanent and unrestricted availability of the progeny of the culture of the deposit to the public upon issuance of the pertinent U.S. patent or upon laying open to the public of any U.S. or foreign patent application, whichever comes first, and assures availability of the progeny to one determined by the U.S. Commissioner of Patents and Trademarks to be entitled thereto according to 35 USC '122 and the Commissioner's rules pursuant thereto (including 37 CFR '1.14 with particular reference to 886 OG 638).

The assignee of the present application has agreed that if a culture of the materials on deposit should die or be lost or destroyed when cultivated under suitable conditions, the materials will be promptly replaced on notification with another of the same. Availability of the deposited material is not to be construed as a license to practice the invention in contravention of the rights granted under the authority of any government in accordance with its patent laws.

The foregoing written description is considered to be sufficient to enable one skilled in the art to practice the invention. The present invention is not to be limited in scope by the example presented herein. Indeed, various modifications of the invention in addition to those shown and described herein will become apparent to those skilled in the art from
5 the foregoing description and fall within the scope of the appended claims.

WHAT IS CLAIMED IS:

1. An isolated WISP-1 antagonist which inhibits or neutralizes induction or secretion of HAS2, HA, CD44 or RHAMM by native WISP-1 polypeptide in at least one type of mammalian cell, said antagonist being selected from the group consisting of an anti-WISP-1 antibody, a WISP-1 immunoadhesin and a WISP-1 variant.
2. The antagonist of claim 1, wherein said antagonist comprises an anti-WISP-1 antibody.
3. The antagonist of claim 2, wherein said anti-WISP-1 antibody binds native human WISP-1 polypeptide comprising amino acids 23-367 of Figures 9A-9C (SEQ ID NO:1) or one or more domains of WISP-1 polypeptide comprising amino acids encoded by the sequences of SEQ ID NO:3, 4, 5, 6, 7, 8, 9, 10 or 11.
4. The antagonist of claim 1, wherein said antagonist comprises a WISP-1 immunoadhesin.
5. A composition comprising the antagonist of Claim 1 and a carrier.
6. The composition of Claim 5 wherein said carrier is a pharmaceutically-acceptable carrier.
7. A method of inhibiting or neutralizing WISP-1 induction or secretion of HAS2, HA, CD44 or RHAMM in mammalian cells, comprising exposing said mammalian cells to an effective amount of WISP-1 antagonist, wherein said WISP-1 antagonist is selected from the group consisting of
 - d) a WISP-1 immunoadhesin;
 - e) a WISP-1 polypeptide linked to a nonproteinaceous polymer selected from the group consisting of polyethylene glycol, polypropylene glycol, and polyoxyalkylene;
 - f) a WISP-1 antibody; and
 - d) a WISP-1 variant.
8. The method of claim 7, wherein said WISP-1 immunoadhesin comprises a WISP-1 sequence fused to a Fc region of an immunoglobulin.
9. The method of claim 7, wherein said anti-WISP-1 antibody binds native human WISP-1 comprising amino acids 23-367 of Figures 9A-9C (SEQ ID NO:1)

or one or more domains of WISP-1 polypeptide comprising amino acids encoded by the sequences of SEQ ID NO:3, 4, 5, 6, 7, 8, 9, 10 or 11.

5 10. The method of claim 7, wherein said mammalian cells comprise cancer cells.

11. The method of claim 10, wherein said mammalian cells comprise colon or colorectal cancer cells, breast cancer cells, lung cancer cells or brain cancer cells.

12. A method of treating cancer in a mammal, comprising administering to said mammal an effective amount of WISP-1 antagonist, wherein said antagonist is selected from the group consisting of an anti-WISP-1 antibody, a WISP-1 immunoadhesin and a WISP-1 variant.

13. The method of claim 12, wherein said cancer comprises colon or colorectal cancer cells, breast cancer cells, lung cancer cells, or brain cancer cells.

14. The method of claim 12, wherein said antagonist inhibits or reduces cancer cell growth or metastasis.

15. The method of claim 12, wherein chemotherapy or radiation is administered to said mammal.

16. The method of claim 12 wherein said antagonist wherein said antagonist inhibits or neutralizes induction or secretion of HAS2, HA, CD44 or RHAMM by native human WISP-1 polypeptide in at least one type of mammalian cell.

17. The method of claim 12 wherein said antagonist is an anti-WISP-1 antibody.

18. The method of claim 17 wherein said anti-WISP-1 antibody binds native human WISP-1 comprising amino acids 23-367 of Figures 9A-9C (SEQ ID NO:1) or one or more domains of WISP-1 polypeptide comprising amino acids encoded by the sequences of SEQ ID NO:3, 4, 5, 6, 7, 8, 9, 10 or 11.

19. A monoclonal antibody comprising the 3D11, 11C2, 9C10, 5D4, or 9C11 antibody secreted by the hybridoma deposited with ATCC as accession number _____, _____, _____, _____, or _____, respectively.

20. A monoclonal antibody which binds to the same epitope as the epitope to which the 3D11, 11C2, 9C10, 5D4, or 9C11 monoclonal antibody produced by the hybridoma cell line deposited as ATCC accession number _____, _____, _____, _____, or _____, respectively, binds.
- 5 21. The hybridoma cell line which produces monoclonal antibody 3D11, 11C2, 9C10, 5D4, or 9C11 produced by the hybridoma cell line deposited as ATCC accession number _____, _____, _____, _____, or _____, respectively.
- 10 22. An isolated anti-WISP-1 antibody, comprising an antibody which binds to WISP-1 polypeptide and competitively inhibits binding of the monoclonal antibody 3D11, 11C2, 9C10, 5D4, or 9C11 produced by the hybridoma cell line deposited as ATCC accession number _____, _____, _____, _____, or _____, respectively, to said WISP-1 polypeptide.
- 15 23. The antibody of claim 20 which is a chimeric, human or humanized antibody.
- 20 24. A chimeric anti-WISP-1 antibody which specifically binds to WISP-1 polypeptide and said antibody comprises a sequence derived from the 3D11, 11C2, 9C10, 5D4, or 9C11 monoclonal antibody produced by the hybridoma cell line deposited as ATCC accession number _____, _____, _____, _____, or _____, respectively.
- 25 25. The antibody of claim 24 wherein said derived sequence is the variable or hypervariable region of the 3D11, 11C2, 9C10, 5D4, or 9C11 monoclonal antibody.





Figure 1




Figure 2




Figure 3




Figure 4




Figure 5

Figure 6






Figura 7



a



b




Figure 8

1 CCCACGGGTC CGCTGGGCCC AGCTCCCCCG AGAGTGGTC GGATCCTCTG GGCTGTCTCG TCGATGCTG TGCCACTGAC GTCCAGGCAT GAGGTGCTTC
GGGTGCGCAG GCGACCCGGG TCGAGGGGGC TCTCCACCAG CCTAGGAGC CCGACAGCC AGCTACGGAC ACGGTGACTG CAGGTCCGTA CTCCACCAAG
M R W F
101 CTGCCCTGGA CGCTGGGAGC AGTGACAGCA GCAGCGGCCA GCACCTCTCT GGCACGGGCC CTCTCTCCAG CCCTACGAC CATGGACTTT ACTCCAGCTC
GAGGGGACCT GGGACCGTGG TCACTGTCTGT CGTGGCGGT CGTGAGGA CCGGTGCCGG GAGAGAGCTC GGGATGCTG GTACTGAAA TGAGGTGGAG
5 L P W T L A A V T A A A A S T V L A T A L S P A P T T M D F T P A P
201 CACTGGAGGA CACCTCTTCA CGCCCCCAAT TCTGCAAGTG GCCATGTGAG TGCCCCCAT CCCACCCCG CTGCCCTGCT GGGTCAGCC TCATCACAGA
GTGACCTCTCT GTGGAGGAGT GCGGGGGTTA AGACTTCTAC CGGTACTACT ACGGGCGGTA GGGGTGGGGC GACGGGCGAC CCCAGTCTGG AGTAGTGTCT
39, L E D T S S R P Q F C K W P C E C P P S P P R C P L G V S L I T D
301 TGCTGTGAG TGCTGTAGA TGTGGCTCA GCAGCTTGG GACAATGCA CGGAGCTGC CATCTGTGAC CCCACCGGG GCCTCTACTG TGACTACAGC
ACCGACACTC ACGACATTCT ACACGGGAGT CGTCGACCG CTGTTGACGT GCCTCGACG GTAGACACTG GGGTGGGCC CGGAGATGAC ACTGATGTCTG
72 G C E C C K M C A Q Q L G D N C T E A A I C D P H R G L Y C D Y S
401 GGGGACGCC CGAGGTACGC AATAGGAGTG TGTGCACAGG TGGTGGTGT GGGCTGGCTC CTGGATGGGG TGGCTACAA CAACGGCCAG TCCTTCCAGC
CCCCGGGG GTCCATGCG TTATCTCTAC ACACGTGTCC ACCAGCACA CCGAGCCAG GACCTACCC ACGGATGTT GTTGGCGGTG AGGAAGGTCTG
105 G D R P R Y A I G V C A Q V V G V G C V L D G V R Y N N G Q S F Q P
501 CTAACCTGAA GTACAACTGC ACGTGCATCG ACGGCGCGGT GGGCTGCACA CCACTGTGCC TCGAGTGGC CCCCCCGGT CTCTGGTGCC CCCACCGCG
GATTGACGTT CATGTTGAGG TGCACGTAGC TGCCCGGCCA CCGACGTGT GGTGACAGG AGSETCACGC GGGGGCGCA GAGACCACGG GGGTGGGCGC
139 N C K Y N C T C I D G A V G C T P L C L R V R P P R L W C P H P R
601 GCGGTGAGC ATACCTGGCC ACTGCTGTA GCAGTGGGTA TGTGAGAGC AGGCCAAGAG GCCACGCAAG ACCGACCCC GTGACACAGG AGCCTTCGAT
CGCGCACTCG TATGGACCGG TGACGACACT CGTCACCCAT ACATCTCTGC TCGGTTCTC CCGTCCGTTT TGGCTGGGG CACTGTGTCC TCGGAAGCTA
172 R V S I P G H C C E Q W V C E D D A K R P R K T A P R D T G A F D
701 GCTGTGGTG AGGTGGAGGC ATGSCACAGG AACTGCATAG CCTACACAAG CCCTGGAGC CCTTGTCTCA CCAGCTGGG CCTGGGGTC TCCACTCGGA
CGACACCCAC TCCACCTCG TACCGTGTCC TTGACGTATC GGATGTGTTT GGGGACCTCG GGAACGAGGT GGTGACAGCC GGACCCCGAG AGGTGAGCCT
205 A V G E V E A W H R N C I A Y T S P W S P C S T S C G L G V S T R I
801 TCTCCAAATGT TAAAGCCCGG TGCTGGCCTG AGCAAGAGAG CCGCTCTGCG AACTTGGGCG CATGGATGT GGACATCCAT ACACATCATTA AGGCAGGGAA
AGAGGTTACA ATTGCGGGTC ACGACCGGAC TCGTTCTCTC GCGGGAGAGG TTGAACGCGG GTACCTACA CCTGTAGTA TGTGAGTAAT TCGTCCCTT
239 S N V N A Q C W P E Q E S R L C N L R P C D V D I H T L I X A G K

FIG.-9A

901 GAAGTGTCTG GCTGTGTACC AGCCAGAGGC ATTCATGAAC TTCCACACTTG CGGCTGCAT CAGCACACCC TCCTATCAAC CCAAGTACTG TGGAGTTTGC
 CTTACACAGC CGACACATGG TCGGTCTCCG TAGGTACTTG AAGTGTGAAC GCCCGACGTA GTGCTGTGGG AGGATAGTTG GGTTCATGAC ACCTCAAAAG
 272 K C L A V Y Q P E A S M N F T L A G C I S T R S Y Q P K Y C G V C
 1001 ATGGACAATA GGTGCTGCAT CCCCTACAAG TCCTAAGACTA TCGACGTCTC CTTCAGTGT CTGATGSGC TTGGCTTCTC CGCCAGGTC CTATGGATTA
 TACCTGTAT CCAGGACGTA GGGGATGTTT AGATTCTGAT AGCTGCACAG GAAGTCCACA GGACTACCG AACCGAAGAG GCGGTCCAG GATACCTAAT
 305 M D N R C C I P Y K S K T I D V S F Q C P D G L G F S R Q V L W I N
 1101 ATGCCCTGCTT CTGTAACTG AGCTGTAGGA ATCCCAATGA CATCTTTGCT GACTTGGAT CCTACCTGA CTCTCAGAA ATTGCCAACT AGCAGGAC
 TACGGACGAA GACATTGGAC TCGACATCTT TAGGGTACT GTAGAAACGA CTGAACCTTA GGATGGACT GAAGCTCTT TAACGGTTGA TCCGTCCGTG
 339 A C F C N L S C R N P N D I F A D L E S Y P D F S E I A N O
 1201 AAATCTTGGG TCTTGGGAC TAACCCCAATG CCGTGAAGC AGTCAGCCCT TATGGCCAT AACTTTTCAC CAATGACCT TAGTTACCCCT GATCTGGACC
 TTTAGAACCC AGAACCCCTG ATTTGGTTAC GGACACTTCG TCAGTCGGA ATACCGGTTA TTGAAGAAGTG GTTACTCGA ATCAATGGGA CTAGACCTGG
 1301 CTTGGCTCC ATTCTGTCT CTAAACCATTC AATGACGCC TGATGTGCT GCTCAGGCC ATGCTATGAG TTTCTCTCTT GATATCATTC AGCATCTACT
 GAACCGGAGG TAAAGACAGA GATTGTAAAG TTTACTCGG ACTACCAGA CGAGTCGGG TACGATACTC AAAAGAGAA CTATAGTAAG TCGTAGATGA
 1401 CTAAAGAAA ATGCTGTCT CTAGCTGTTT TGGACTACAC CCAAGCTGA TCCAGCCTTT CCAAGTCACT AGAAGTCTTG CTGGATCTTG CCTAAATCCC
 GATTTCTTTT TACGGACAGA GATCGACAG ACCTGATGTT GGTTCGACT AGTTCGAAA GGTTCACTGA TCTTCAGGAC GACCTAGAAC GGATTTAGGG
 1501 AAGAAATGGA ATCAGGTAGA CTTTAAATAT CACTAATTC TTCTTTAGAT GCCAAACCAC AAGACTTTT GGGTCCATTC AGATGAATAG ATGGAATTTG
 TTCTTTACCT TAGTCCATCT GAAATTTATA GTGATTTAAG AAGAAATCTA CGGTTTGGTG TTCTGAGAA CCCAGGTAAG TCTACTTATC TACCTTAAAC
 1601 GAACAATAGA ATAATCTATT ATTTGGAGCC TGGCAGAGG TACTGTAAAG GGTAAATCTG AGTCAAGCC ACCAAACTA TCCTGATTC AAATATGTAT
 CTTGTATCT TATTAGATA TAAACCTCGG ACGTTCTCC ATGACATTAC CCATTAAAG TGCAGTCCG TGGTTTGTAT AGGACTAAGG TTTATACATA
 1701 GCACCTCAAG GTCATCAAC ATTTGCCAAG TGAGTTGAAT AGTTGCTTAA TTTTGATTTT TAATGGAAG TTGTATCCAT TAACCTGGGC ATTTGTGAGG
 CGTGGAGTTC CAGTAGTTTG TAAACGGTTC ACTCAACTTA TCACGAAT AAAACTAAAA ATTACCTTTC AACATAGTA ATTTGACCCG TAACAATCC
 1801 TTAAGTTTCT CTTACCCCTT ACACGTGMA GGGTACAGAT TAGTTTGTG CCAGTCAGAA ATAAATTTG ATAAACATTC CTGTTGATGG GAAAGCCCC
 AATTCAAAAGA GAGTGGGGA TGTGACACTT CCCATGTCTA ATCCAAACAG GGTCACTCTT TATTTTAAAC TATTTGTAAG GACAACCTACC CTTTTCGGG
 1901 CAGTTAATAC TCCAGAGACA GGAAGGTC AGCCATTTT AGAAGGACCA ATTGACTCTC ACACGTATC AGCTGTGAC TGGCAGGGCT TTGGGCAATT
 GTCAATTATG AGGTCTCTGT CCTTTCCAG TCGGTAAAG TCTTCTGTT TAACGTAGAG TGTGACTAG ACCGTCCGA AACCCGTCAA

FIG.-9B

2001 GGCACGGCTC TTCCTTGAAT CTCTCTCCCTT GTCTCTGCTTG GGTTCATAGG AATTGGTAAG GCCTCTGGAC TGGCCTGTCT GGCCTCTGAG AGTGGTGCCC
 CCGTCCGAG AAGGAACCTTA GAAGAGGGAA CAGGACGAAC CCAAGTATCC TTAACCATTC CGAGAGCCTG ACCGGGACTC CCGGGACGGG TCACACCGGG
 2101 TGGAAACACTC CTCTACTCTT ACAGAGCCCTT GAGAGACCCA GGTGCAGACC ATGCCAGACC CACTGAATG ACCAAGACAG GTTCAGGTAG GGTGTGGGT
 ACCTGTGTGAG GAGATGAGAA TGCTCTGGAA CTCTCTGGGT CGACGTCTGG TAGCGTCTGG GTGACTTTTAC TGGTCTGTGC CAAGTCCATC CCCACACCCA
 2201 CAACCAAGA AGTGGGTGCC CTTCGTAGCA GCCTGGGGTG ACCTCTAGAG CTGGAGGCTG TGGACTCCA GGGGCCCTCG TGTTCAGGAC ACATCTATTG
 GTTGGTTCT TCACCCACGG GAACCATCGT CGGACCCAC TGGAGATCTC GACCTCCGAC ACCCTGAGGT CCGCGGGGCG ACAAGTCTTG TGTAGATAAC
 2301 CAGAGACTCA TTTCACAGCC TTTCGTTCTG CTGACCAAAAT GGCAGTTTT CTGGTAGGAA GATGGAGSTT TACCAGTTGT TTAGAAACAG AAATAGACTT
 GTCTCTGAGT AAAGTGTGG AAAGCAAGAC GACTGGTTTA CCGGTCAAAA GACCATCCTT CTACCTCCA ATGGTCAACA ATCTTTGTCT TTATCTGAA
 2401 AATAAAGGT TAAAGCTGAA GAGGTGAA CTAAAGGAA AAGGTGTTG TTAATGAATA TCAGGCTATT ATTTATTGTA TTAGGAATAT ATAATATTTA
 TTATTTCCAA ATTTCGACTT CTCCAACCTC GATTTTCCTT TTCCAACAAC AATTACTTAT AGTCCGATAA TAAATTAACAT AATCTTTTA TATTATAAAT
 2501 CTGTTAGAA TCTTTTATTT AGGGCCTTTT CTGTGCCAGA CATGTCTCTC AGTGCTTTGC ATGTATTAGC TCACTGAATC TTCACGACAA TGTGAGAG
 GACAATCTTA AGAAATATAA TCCCGGAAAA GACACGGTCT GTAACGAGAG TCACGAAACG TACATAATCG AGTGACTTAG AAGTGTGT ACAACTCTTC
 2601 TTCCCATTTAT TATTTCTCTT CTACAAATG TGAACGGAA GCTCATAGAG GTGAGAAAC TCACCCAGAG TCACCCAGTT GGTGACTGGG AAAGTTAGGA
 AAGGTAATA ATAAAGACAA GAATGTTTAC ACTTTGCCCTT CGAGTATCTC CACTCTTTTG AGTGGTCTC AGTGGGTCAA CCACTGACCC TTTCATCTCT
 2701 TTCAGATCGA AATTGGACTG TCTTTATNAC CCATATTTTC CCCCTGTTTT TAGAGCTTCC AAATGTGTCA GAATAGGAAA ACATTGCAAT AAATGGCTTG
 AAGTCTAGCT TTAACCTGAC AGAATATTG GGTATAAAG GGGGACAAA ATCTCGAAGG TTTACACAGT CTTATCTCTT TGTAACTTA TTTACCGAAC
 2801 ATTTTAAAA AAAAAAATAA AAAAAAATAA
 TAAAAAATTT TTTTTTTTTT TTTTTTTTTT

FIG.-9C

Effect of WISP-1 expression on NRK metastatic potential				
No. Cells injected	Weeks after inoculation	Incidence of lung metastasis		
		NRK/control	NRK/WISP-1L	NRK/WISP-1H
2.5×10^5	2	0/1	0/1	2/2 (grade I)
	3	0/4	3/3 (grade I)	4/4 (grade I-II)
	4	0/4	4/4 (grade I)	3/3 (grade III)
0.5×10^5	2	0/2	0/2	1/2 (grade I)
	3	0/4	2/4 (grade I)	3/4 (grade I)
	4	0/4	4/4 (grade I)	3/4 (grade I-II)

Fig. 10




Figure 5




Figure 6





Figure 7

a



b




Figure 8

1 CCCACGGTC CGCTGGGCC AGCTCCCCG AGAGGTGGTC GGATCCTCTG GGTGCTCGG TCGNTGCTG TCCACTGAC GTCCAGGCAT GAGTGGTTC
GGGTGGCAG GCGACCCGG TCGAGGGGC TCCTCCACCAG CCTAGGAGAC CCGACGACC AGCTACGGAC AGGTGACTG CAGTCCGTA CTCACCAAG
M R W F
1
101 CTGCCCTGGA CGCTGGCAGC AGTGACAGCA GCAGCCGCCA GCACGTCTCT GGCACGGCC CTCTCTCCAG CCCCTACAC CATGGACTTT ACTCCAGTCT
GACGGACCTT CCGACCCGTC TCACGTGTCGT CGTGGCCGGT CGTGCGAGGA CCGGTGCCG GAGAGAGTCT GGGATGCTG GTACCTGAAA TGAGTCCGAG
5 L P W T L A A V T A A A A S T V L A T A L S P A P T T M D F T P A P
201 CACTGGAGGA CACCTCTCTA CGCCCCCAAT TCTGCAAGTG GCCATGTAG TCCCGCCAT CCCACCCCG CTGCCCGCTG GGGTCAGCC TCATCAGCA
GTACCTCTT GTGGAGGAGT GCGGGGGTTA AGACCTTCAC CGGTACACTC ACGGGGGTA GGGGTGGGC GACGGGCGAC CCCAGTCCG AGTAGTGTCT
39. L E D T S S R P Q P C K W P C E C P P S P P R C P L G V S L I T D
301 TGCGTGTAG TGCTGTAGA TGTGCGCTCA GCAGTTTGG GACAATGCA CGAGGCTGC CATCTGTGAC CCCACCGGG GCCTCTACTG TGAATACAGC
ACGACACTC ACGACATTT ACACCGAGT CGTCCGAGT CGTCCGAGT GCTCTCGACG GTAGACACTG GGGTGGGCC CGGAGATGAC ACTGATGTCTG
72 G C E C C K M C A Q Q L G D N C T E A A I C D P H R G L Y C D Y S
401 GGGGACCCG CAGGTACGC AATAGGAGTG TGTGACAGG TGGTGGTGT GGGTGGTCT CCGGTGGGG TCGGTACAA CAACGGCCAG TCCTTCAGC
CCCTGGGGG GCTCCATGG TTATCTCTAC ACAGTGTTC ACCAGCACA CCGACGCGAG GACTACCCC ACGGATGTT GTTGGCGTC AGGAAGTCTG
105 G D R P R Y A I G V C A Q V V G V G C V L D G V R Y N N G Q S F Q P
501 CTAACTGCAA GTACAACTGC ACGTGCATCG ACGGGCGGT GGGCTGCACA CCACTGTGCT TCCAGTGGC CCCCGCGGT CTCTGGTCC CCCACCCGG
GATTGACGTT CATGTTGACG TGCACGTAGC TCGCGGCCA CCGGACGTT GTGACACGG AGGTCACGC GGGGGCGCA GAGACCACGG GGTGGGGC
139 N C K Y N C T C I D G A V G C T P L C L R V R P P R L W C P H P R
601 GCGGTGAGC ATACTGGCC ACTGTGTGA GCAGTGGTA TGTGAGGAG ACGCAAGAG GCCAGCAAG ACCGACCCC GTGACACAGG AGCTTCGAT
CGGCACTCG TATGGACCG TGACGACACT CCGTACCCAT ACATCTCTG TCGGTCTC CGGTGGTCT TGGCTGGGG CACTGTGTC TCGGAGCTA
172 R V S I P G H C C E Q W V C E D D A K R P R K T A P R D T G A F D
701 CTTGGGTG AGTGGAGC ATGGACAGG AACTGCATAG CCTACACAAG CCCCTGGAG CCCCTGCCA CCACTGGGG CTTGGGGTC TCCACTCGGA
CGACACCCAC TCCACCTCG TACCGTGTCC TTGACGTATC GGATGTTC GGGACCTCG GGAACGAGT GTGCGACGC GGAACCCCG AGGTAGCCT
205 A V G E V E A W H R N C I A Y T S P W S P C S T S C G L G V S T R I
801 TCTCCAATGT TAACGCCAG TGTGGCTG AGCAAGAG CCGCTCTGC AACTTGGGC CATGCGATGT GGAATCCAT ACATCATTA AGGCAGGAA
AGAGGTTACA ATTGGGCTC ACGACCGAC TCGTCTCTC GCGGAGACG TTGAACGCG GTACGCTACA CCTGTAGTA TGTAGTAT TCCGTCTT
239 S N V N A Q C W P E Q E S R L C N L R P C D V D I H T L I K A G K

FIG.-9A

901 GAAGTGTCTG GCTGTGTACC AGCCAGAGGC ATCCATGAAC TTCACACTTG CGGGTGCAT CAGCACACGC TCCTATCAAC CCAAGTACTG TGGAGTTTGC
 CTTACACAGC CGACACATGG TCGGTCTCCG TAGGTACTTG AAGTGTGAAC GCCCAGCGTA GTCGTGTGCG AGGATAGTTG GGTTCATGAC ACCTCAACAG
 272 K C L A V Y Q P E A S M N F T L A G C I S T R S Y Q P K Y C G V C
 1001 ATGGACAATA GGTGCTGCAT CCCATACAAG TCTAAGACTA TCGAGGTGTC CTTCCAGTGT CTTGATGGGC TTGGGTTCTC CGCCAGGTC CTATGGATTA
 TACCTGTTAT CCACGACGTA GGGGATGTTT AGATTCTGAT AGCTGCACAG GAAGTCCACA GGACTACCCG AACCGAAGAG GCGGTGCCAG GATACCTAAT
 305 M D N R C C I P Y K S K T I D V S F Q C P D G L G F S R Q V L W I N
 1101 ATGCTCTGCT CTGTAACCTG AGCTGTAGGA ATCCCAATGA CATCTTTGCT GACTTGGAT CTTACCCCTGA CTTCTCAGAA ATTGCCAACT AGGCAGGCAC
 TACGGACGAA GACATTTGAC TCGACATCCT TAGGGTTACT GTAGAAACGA CTGAACCTTA GGATGGGACT GAAGGTCTT TAAACGGTTGA TCCGTCCGTC
 339 A C F C N L S C R N P N D I F A D L E S Y P D F S E I A N O
 1201 AATCTTTGG TCTTGGGAC TAACCCAAATG CTTGTGAAGC AGTCAGCCCT TATGGCCAAAT AACTTTTTC CATTAGCCT TAGTTACCCCT GATCTGGACC
 TTAGAAACCC AGAACCCCTG ATTGGGTTAC GGACACTTCG TCAGTCCGGA ATACCCGTTA TTGAAAGCTG GTTACTCGA ATCAATGGGA CTAGACCTGG
 1301 CTTGGCCTCC ATTCTCTCT CTAACCAATC AATGAGGCC TGATGTGCT GCTCAGGCC GCTCAGGCC AACTTTTTC CATTAGCCT TAGTTACCCCT GATCTGGACC
 GAACCGGAGG TAAAGACAGA GATTGGTAAG TTACTCTGG ACTACCAGGA CGAGTCCGG TACCATCTC TAAAGAGAA CTATAGTAAG TCGTAGATGA
 1401 CTAAGAGAAA ATGCTGTCT CTAGCTGTC TGGACTACAC CCAAGCCTGA TCCAGCCCTT CCAAGTCACT AGAAGTCTG CTGGATCTTG CCTAAATCCC
 GATTTCTTTT TACGGACAGA GATCGACAG ACCTGATGTG GGTTCGACT AGGTCGAAA GGTTCAGTGA TCTTCAGGAC GACCTAGAAC GGATTTAGGG
 1501 AAGAAATGGA ATCAGGTAGA CTTTAAATAT CACTAATTC CACTAATTC TTCTTAGAT GCCAAACCAC AAGACTCTTT GGTCTCATTC AGATGAATAG ATGGAATTTG
 TTCTTTACCT TAGTCCATCT GAAATATATA GTGATTAAAG AAGAAATCTA CGGTTTGGTG TTCTGAGAA CCCAGSTAAG TCTACTTATC TACCTTAAAC
 1601 GAACAATAGA ATATCTATT ATTGGAGCC TGGCAGAGG TACTGTATG GGTAAATCTG AGGTACGCG AGGTACGCG ACCAAACTA TCCTGATTC AAATATGTAT
 CTTGTATCT TATTAGATAA TAAACCTCG ACGTTCTCC ATGACATTAC CCATTAGAC TGCAGTCCG TGGTTTGTAT AGGACTAAGG TTTATACATA
 1701 GCACCTCAAG GTCATCAAAC ATTTGCCAAG TGAGTTGAT AGTTGCTTAA TTTTGATTTT TAATGGAAG TTGTATCCAT TAACCTGGGC ATTTGTTGAGG
 CGTGAGTTC CAGTAGTTG TAAACGGTTC ACTCAACTTA TCAAGGAAT TAAACTAAAA ATTACCTTC AACATAGTA ATTGGACCCG TAACAACCTC
 1801 TTAAGTTTCT CTTACCCCT ACACGTGAA GGGTACAGAT TAGGTTTGT CCACTCAGAA ATAAATTTG ATAAACATTC CTGTTGATGG GAAAGCCCC
 AATCAAAGA GAAGTGGGA TGTGACACTT CCCATGCTA ATCCAAACAG GGTCACTCT TATTTTAAAC TATTTGTAAG GACAACATACC CTTTTCGGGG
 1901 CAGTAAATAC TCCAGAGACA GGGAAAGGTC AGCCCATTC AGAAGGACCA ATTGACTCT ACCTGAATC AGCTCTGAC TGGCAGGCT TTGGGCAAT
 GTCAATATG AGGTCTCTGT CCTTTCCAG TCGGTAAG TCTTCTGAT TAACTGAG TGTGACTTAG TCGAGACTG ACCGTCCCG AACCCGTCAA

FIG.- 9B


2001 GGCACGGCTC TTCTTGAAT CTCTCCCTT GTCTGCTTG GGTTCATAGG AATTGTAAAG GCCTCTGGAC TGGCCTGTCT GGCCTCTGAG AGTGTGCCCC
 CCGGTCCGAG AAGGAACCTTA GAAGAGGGA CAGGACGAA CCAAGTATCC TTAACCATTC CGGAGACCTG ACCGGACAGA CCGGGACTC TCACCACGGG
 2101 TGGAACACTC CTCTACTCTT ACAGAGCCTT GAGAGACCCA GCTGCAGACC ATGCCAGACC CACTGAATG ACCAAGACAG GTTCAGGTAG GGGTGTGGGT
 ACCTTGTGAG GAGATGAGAA TGCTCTGGAA CTCTCTGGGT CGACTCTGG TACGGTCTGG GTGACTTTAC TGGTCTCTGC CAAGTCCATC CCCACACCCA
 2201 CAAACCAAGA AGTGGGTGCC CTGGGTAGCA GCCTGGGTG ACCTCTAGAG CTGGAGGCTG TGGGACTCCA GGGGCCCCCG TGTTCAGGAC ACATCTATTG
 GTTGTGTTCT TCACCCACGG GAACCATCGT CGGACCCAC TGGAGATCTC GACCTCCGAC ACCCTGAGGT CCGCGGGGC ACAAGTCCCTG TGTAGATAC
 2301 CAGAGACTCA TTTCACAGCC TTTCGTTCTG CTGACCAAT GGCAGTTTT CTGGTAGGAA GATGGAGTT TACCAGTTGT TTAGAACAG AAATAGACTT
 GTCTCTGAGT AAGTGTGG AAAGCAAGAC GACTGTGTTA CCGTCAAAA GACCATCTT CTACCTCAA ATGCTCAACA AATCTTGT TTTATCTGAA
 2401 AATAAAGGT TAAAGCTGAA GAGTTTGAAG CTAAAGGAA AAGTTGTGTT TTAATGAATA TCAGGCTATT ATTTATTGTA TTAGGAAAT ATAAATTTTA
 TTATTTCCAA ATTTGACTT CTCCAACCTC GATTTTCCCT TTCCAACNAC AATTACTTAT AGTCCGATAA TAAATAACAT AATCCTTTTA TATTATAAT
 2501 CTGTAGAAAT TCTTTTATTT AGGCCCTTTT CTGTGCCAGA CATTGCTCTC AGTCTTTGC ATGTATTAGC TCACTGAATC TTCACGACAA TGTGAGAAG
 GACAATCTTA AGAAATATAA TCCCGGAAA GACACGGTCT GTAACGAGAG TCACGAAACG TACATATCG AGTGACTTAG AAGTGTGTT ACACTCTTC
 2601 TTCCCATAT TATTCTGTGTT CTTACAATG TGAACGGAA GCTCATAGAG GTGAGAAAC TCAACCAAG TCACTCCAGTT GGTGACTGGG AAAGTTAGGA
 AAGGTAATA ATAAAGACAA GAATGTTTAC ACTTGTGCTT CGAGTATCTC CACTCTTTTG AGTGTGCTC AGTGGGTCAA CCACTGACCC TTTCATCTCT
 2701 TTCAGATGA AATTGACTG TCTTTATAAC CCATATTTTC CCCCTGTTTT TAGAGCTTCC AATGTGTCA GAATAGGAA ACATTCGAAT AAATGCTTG
 AAGTCTAGCT TTAACCTGAC AGAATATTG GGTATAAAG GGGACAAAA ATCTCGAAGG TTTACACAGT CTTATCTCTT TGTACGTTA TTTACCGAAC
 2801 ATTTTATAA AAAAAAAA AAAAAAAA
 TAAAAAATTT TTTTTTTT TTTTTTTT

FIG.-9C


5' 400 3' 1000 2000 3000 4000 5000

Effect of WISP-1 expression on NRK metastatic potential				
No. Cells injected	Weeks after inoculation	Incidence of lung metastasis		
		NRK/control	NRK/WISP-1L	NRK/WISP-1H
2.5×10^5	2	0/1	0/1	2/2 (grade I)
	3	0/4	3/3 (grade I)	4/4 (grade I-II)
	4	0/4	4/4 (grade I)	3/3 (grade III)
0.5×10^5	2	0/2	0/2	1/2 (grade I)
	3	0/4	2/4 (grade I)	3/4 (grade I)
	4	0/4	4/4 (grade I)	3/4 (grade I-II)

Fig. 10



WISP-1: Wnt-1 Induced Secreted Protein



Deletion Mutants of Human WISP-1

Name IGFBP VWC variable TSP CT

1,2,3,4

1,2,3

2,3,4

1,2,4

1,3,4

1,2


1

2

3


4

Epitope of WISP-1 Antibodies clones 11C2, 5D4 and 9C11



The epitope of the WISP-1 antibodies clones 11C2, 5D4 and 9C11 is located within the first 19 amino acids of WISP-1 variable region.

Epitope Mapping of WISP-1 Antibody 3D-11



▲ The epitope of the WISP-1 antibody 3D11 is located within WISP-1 domain 1.





Figure 12

Epitope Mapping of Anti-WISP-1 Mab (11C2)

Name	IGFBP	VWC	variable	TSP	CT	WB
1,2,3,4						+
1,2,3						+
2,3,4						+
1,2,4						-
1,3,4						-
1,2						-
1						-
2						-
3						-
4						-




Figure 13




Figure 14










Figure 15

WISP-1 Antibodies Characterization

Clones	Isotype	Epitope		IP	WB	Blocking Activity		
		Struct.	Cross Compet.			Heparin Binding	Cell Migrat.	Lung Met.
3D11	2b	Domain 1	A	-	+	+/-	+	+
9C10	1	Domain 1	B	+	-	-	+	+
11C2	2b	Variable region	C	+	+	+	+	+
5D4	2a	Variable region	C	+	+	+	+	+
9C11	2a	Variable region	C	+	+	+	+	+



United States Patent & Trademark Office
Office of Initial Patent Examination -- Scanning Division



Application deficiencies found during scanning:

☐ Page(s) _____ of _____ were not present
for scanning. (Document title)

☐ Page(s) _____ of _____ were not present
for scanning. (Document title)

Scanned copy is best available. Most of the drawings are very dark

**This Page is Inserted by IFW Indexing and Scanning
Operations and is not part of the Official Record**

BEST AVAILABLE IMAGES

Defective images within this document are accurate representations of the original documents submitted by the applicant.

Defects in the images include but are not limited to the items checked:

☐ BLACK BORDERS

☐ IMAGE CUT OFF AT TOP, BOTTOM OR SIDES

☐ FADED TEXT OR DRAWING

☐ BLURRED OR ILLEGIBLE TEXT OR DRAWING

☐ SKEWED/SLANTED IMAGES

☒ COLOR OR BLACK AND WHITE PHOTOGRAPHS

☐ GRAY SCALE DOCUMENTS

☒ LINES OR MARKS ON ORIGINAL DOCUMENT

☐ REFERENCE(S) OR EXHIBIT(S) SUBMITTED ARE POOR QUALITY

☒ OTHER: small text

IMAGES ARE BEST AVAILABLE COPY.

As rescanning these documents will not correct the image problems checked, please do not report these problems to the IFW Image Problem Mailbox.